Martin Goßner · Ulrich Ammer

The effects of Douglas-fir on tree-specific arthropod communities in mixed species stands with European beech and Norway spruce

Published online: 28 March 2006
© Springer-Verlag 2006

Eur J Forest Res (2006) DOI:10.1007/s10342-006-0113-y

In Table 5 of this paper the animal genus designations should have been written out in full instead of abbreviated. The corrected Table 5 is shown below

Table 5 Code of species of the ordination diagrams

Coleoptera	1	<i>Absidia rufotestacea</i> (Letzner, 1845)	Heteroptera	23	<i>Atractotomus magnicornis</i> (Fallén, 1907)
	2	<i>Absidia schoenherri</i> (Dejean, 1837)		24	<i>Cremnocephalus alpestris</i> Wagner, 1941
	3	<i>Adalia decempunctata</i> (Linnaeus, 1758)		25	<i>Deraeocoris lutescens</i> (Schilling, 1837)
	4	<i>Anaspis rufilabris</i> (Gyllenhal, 1827)		26	<i>Orius minutus</i> (Linnaeus, 1758)
	5	<i>Anatis ocellata</i> (Linnaeus, 1758)		27	<i>Parapsallus vitellinus</i> (Scholtz, 1846)
	6	<i>Athous subfuscus</i> (Müller, 1764)	Neuroptera	28	<i>Psallus varians</i> (Herrich-Schaeffer, 1841)
	7	<i>Athous vittatus</i> (Fabricius, 1792)		29	<i>Coniopteryx pygmaea</i> Enderlein, 1906
	8	<i>Brachytarsus nebulosus</i> (Forster, 1771)		30	<i>Hemerobius fenestratus</i> Tjeder, 1932
	9	<i>Clytus lama</i> Mulsant, 1847		31	<i>Hemerobius micans</i> Olivier, 1792
	10	<i>Corticaria abietorum</i> Motschulsky, 1867		32	<i>Hemerobius pini</i> Stephens, 1836
	11	<i>Corticarina lambiana</i> (Sharp, 1910)		33	<i>Parasemidalis fuscipennis</i> (Reuter, 1894)
	12	<i>Cryphalus abietis</i> (Ratzeburg, 1837)		34	<i>Peyerimhoffina gracilis</i> (Schneider, 1851)
	13	<i>Cychramus luteus</i> (Fabricius, 1787)	Araneae	35	<i>Anyphaena accentuata</i> (Walckenaer, 1802)
	14	<i>Ernobius abietis</i> (Fabricius, 1792)		36	<i>Coelotes terrestris</i> (Wider, 1834)
	15	<i>Exochomus quadripustulatus</i> (Linnaeus, 1758)		37	<i>Cryphoea silvicola</i> (C.L. Koch, 1834)
	16	<i>Leptophloeus alternans</i> (Erichson, 1845)		38	<i>Dendryphantes rudis</i> (Sundevall, 1833)
	17	<i>Magdalis nitida</i> (Gyllenhal, 1827)		39	<i>Dismodicus elevatus</i> (C.L. Koch, 1838)
	18	<i>Melanotus castanipes</i> (Paykull, 1800)		40	<i>Drapetisca socialis</i> (Sundevall, 1833)
	19	<i>Pityogenes chalcographus</i> (Linnaeus, 1761)		41	<i>Hahnia pusilla</i> C.L. Koch, 1841
	20	<i>Pityophthorus exsculptus</i> (Ratzeburg, 1837)		42	<i>Lathys humilis</i> (Blackwall, 1855)
	21	<i>Pityophthorus pityographus</i> (Ratzeburg, 1837)		43	<i>Pelecopsis elongata</i> (Wider, 1834)
	22	<i>Polydrusus pallidus</i> (Gyllenhal, 1834)		44	<i>Segestria senoculata</i> (Linnaeus, 1758)

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10342-006-0113-y>

M. Goßner · U. Ammer
Department of Ecosystem and Landscape Management,
Technical University of Munich-Weihenstephan,
Am Hochanger 13, 85354 Freising, Germany

M. Goßner (✉)
Loricula, Schussenstraße 12, 88273 Fronreute, Germany
E-mail: martin.gossner@loricula.de
Tel.: +49-7502-913319
Fax: +49-8161-714671