Nutrients return from leaves and litterfall in a mediterranean cork oak (*Quercus suber* L.) forest in southwestern Spain

Enrique Andivia · Manuel Fernández ·
Javier Vázquez-Piqué · Aranzazu González-Pérez ·
Raúl Tapias

Published online: 21 August 2009
© Springer-Verlag 2009

Erratum to: Eur J Forest Res DOI 10.1007/s10342-009-0274-6

There were errors in the Results section on the third page of the original publication, both in the text and in Table 2. In the first paragraph of the Results section '(10.90% of total litterfall)' should read '(55.82% of total litterfall)' and 'acorn production (2,113.68 g m⁻²)' should read 'acorn production (87.10 g m⁻²)'. Table 2 below shows the correct values.

The errors were due to a miscalculation and do not affect any aspect of the discussion and conclusions of the work.

Table 2 Annual values for different litterfall components in g m⁻² (standard error) and percentage of total litterfall of the year

Year	Twigs	Leaves	Catkins	Acorns	Miscellaneous	Total
March 2004–February 2005	70.12 (24.44) 14.17%	274.40 (55.52) 55.82%	24.24 (13.92) 4.93%	87.10 (64.97) 17.82%	35.64 (7.92) 7.26%	491.50
March 2005–February 2006	49.88 (18.84) 18.32%	189.56 (68.12) 69.60%	14.68 (15.56) 5.39%	2.68 (3.12) 0.98%	15.56 (6.84) 5.71%	272.36
Average	60.00 15.71%	231.96 60.74%	19.44 5.09%	44.89 11.76%	25.60 6.70%	381.89

The online version of the original article can be found under
doi: [10.1007/s10342-009-0274-6](https://doi.org/10.1007/s10342-009-0274-6).

E. Andivia (✉) · M. Fernández · J. Vázquez-Piqué ·
A. González-Pérez · R. Tapias
Department of Forest Sciences, University of Huelva,
Ctra. Palos–La Rábida, s/n. 21819 Palos de la Frontera,
Huelva, Spain
e-mail: enrique.andivia@dcaf.uhu.es