Dis Colon Rectum, Vol.
48, No. 2, February 2005,
pp. 237–242 (DOI:
10.1007/s10350-004-0768-8)

In the article “Long-Term Treatment of High Intestinal Output Syndrome With Budesonide in Patients With Crohn’s Disease and Ileostomy,” by Karl W. Ecker, Andreas Stallmach, Jürgen Löffler, Roland Greinwald and Ulrike Achenbach, on page 239 in the paragraph

Dis Colon Rectum 2005; 48: 1834
DOI: 10.1007/s10350-005-0054-4
© The American Society of Colon and Rectal Surgeons
Published online: 3 May 2005

“Ileostomy Output,” the text from line 20 to line 4 in the next column should read:

“After withdrawal of budesonide, 18 of 22 patients showed an increase of ileostomy output and a decrease after re-exposure to budesonide. In 2 patients, the output increased, but did not decrease at re-exposure, and in 2 patients the output decreased at re-exposure of budesonide without an increase at withdrawal before.”

instead of:

“After withdrawal of budesonide, 20 of 22 patients showed an increase of ileostomy output; after re-exposure, in 20 patients the output decreased again. In 18 patients, the output increased after stopping budesonide treatment and decreased after re-exposure.”

The authors regret the error.