Erratum to: First experiences with a new surgical approach in adult full-thickness burns: single step reconstruction of epidermal, dermal and subcutaneous defects by use of split-thickness skin grafting, a dermal collagen matrix and autologous fat-transfer

M. Keck · H. F. Selig · J. Kober · D. B. Lumenta · H. Schachner · A. Gugerell · L.-P. Kamolz

Published online: 17 August 2013
© Springer-Verlag Wien 2013

Erratum to: DOI 10.1007/s10353-013-0211-2

The original version of this article unfortunately contained mistakes:

Fig. 1e was missing. The correct version of Fig. 1a–e is given on the right.

Reference 14 was listed incorrectly. The corrected reference called:

14. Kamolz L-P, Lumenta DB, Kitzinger HB, Frey M. Tissue engineering for cutaneous wounds: an overview of current standards and possibilities. Eur Surg. 2008;40(1):19–26.

The online version of the original article can be found under doi: <http://dx.doi.org/10.1007/s10353-013-0211-2>

M. Keck, MD (✉) · H. F. Selig, MD · J. Kober, MSc ·
D. B. Lumenta, MD · A. Gugerell, PhD ·
L.-P. Kamolz, MD, PhD, MSc

Vienna Burn Centre, Division of Plastic and Reconstructive
Surgery, Department of Surgery, Medical University of Vienna,
Waehringer Guertel 18–20, 1090 Vienna, Austria
e-mail: maike.keck@meduniwien.ac.at

H. F. Selig, MD
Hospital for Hand Surgery, Rhoen Klinikum AG,
Bad Neustadt/Saale, Germany

D. B. Lumenta, MD · L.-P. Kamolz, MD, PhD, MSc
Division of Plastic, Aesthetic and Reconstructive Surgery,
Department of Surgery, Medical University of Graz, Graz, Austria

H. Schachner
Department of Pathology, Medical University of Vienna,
Vienna, Austria