ERRATUM

Effects of Reference Axes Used During Measurements of Ocular and Corneal Higher-Order Aberrations in Patients Following LASIK

Atsuko Takehara¹, Naoyuki Maeda¹, Sayuri Ninomiya², Takashi Fujikado²,
Yoko Hirohara³, and Toshifumi Mihashi³

¹Department of Ophthalmology, Osaka University Medical School, Suita, Osaka, Japan;

²Department of Applied Visual Science, Osaka University Medical School, Suita, Osaka, Japan;

³Topcon Corporation, Tokyo, Japan

Japanese Journal of Ophthalmology 2006;50:318–322

The correct name of the fifth author should be given as
Yoko Hirohara, not Yoko Hiroha.

Received: July 1, 2005 / Accepted: November 2, 2005

Correspondence and reprint requests to: Naoyuki Maeda, Room E7,
Department of Ophthalmology, Osaka University Medical School,
Yamadaoka 2-2, Suita, 565-0871, Japan
e-mail: nmaeda@ophthal.med.osaka-u.ac.jp

Presented in part as a poster at the Association of Research for Vision
and Ophthalmology annual meeting, Fort Lauderdale, Florida, May 6,
2003. Mr. Mihashi and Ms. Hirohara are employees of Topcon Corpo-
ration. No other author or family member has any proprietary or finan-
cial interest in the instrument used in the study.