CORRECTION

Publisher Correction to: Effect of glaucoma implant surgery on intraocular pressure reduction, flare count, anterior chamber depth, and corneal endothelium in primary open-angle glaucoma

Kyoko Ishida^{1,2} · Naoya Moroto² · Kazuhiro Murata² · Tetsuya Yamamoto²

Published online: 12 March 2018
 © Japanese Ophthalmological Society 2018

Correction to: Jpn J Ophthalmol (2017) 61:334–346
<https://doi.org/10.1007/s10384-017-0512-2>

The publisher regrets that the original version of this article unfortunately contained numerous mistakes that occurred during typesetting. Those errors and their corrections are shown below.

Page	Column	Line	In the original article	Should be
334	1	15	IOP $\geq$ 5 mmHg	IOP $\geq$ 5 mmHg
336	1	17	an IOP $\geq$ 5 mmHg	an IOP $\geq$ 5 mmHg
	1	19	an IOP $\geq$ 5 mmHg	an IOP $\geq$ 5 mmHg
	1	28	IOP < 5 mmHg.	IOP < 5 mmHg.
	1	42	the Chi-square	the chi-square
	2	7	patient's	patients'
338	1	1	IOPs > 18 mmHg	IOPs > 18 mmHg
	1	7	IOPs > 15 mmHg	IOPs > 15 mmHg
	1	8	reduction > 20%	reduction > 20%
	2	5	of < 0.3333	of < 0.3333
341	1	20	IOP $\geq$ 5 mmHg	IOP $\geq$ 5 mmHg
	2	18	IOP $\leq$ 10 mmHg	IOP $\leq$ 10 mmHg
	2	25	IOP $\leq$ 10 mmHg	IOP $\leq$ 10 mmHg
343	1	8	1% to 52.4%	1 to 52.4%
	1	20	Chiou et al	Chiou et al.
	2	42	IOP < 5 mmHg	IOP < 5 mmHg

Page	Column	Line	In the original article	Should be
344	2	33	Conflicts of interest K. Ishida, N. Moroto, K. Murata have no conflict of interest. T. Yamamoto, Grants (Alcon Japan, MSD Japan, Otsuka, Pfizer Japan, Santen, Senju), Consultant/Advisor fees (Alcon Japan, Kowa, Otsuka, Pfizer Japan, Rohto, Senju), Lecture fees (Alcon Japan, AMO Japan, CREWT Medical Systems, Johnson & Johnson, Kowa, MSD Japan, Novartis Pharma, Otsuka, Pfizer Japan, R-Tech Ueno, Santen, Senju).	Conflicts of interest K. Ishida, None; N. Moroto, None; K. Murata, None; T. Yamamoto, Grants (Alcon Japan, MSD Japan, Otsuka, Pfizer Japan, Santen, Senju), Consultant/Advisor fees (Alcon Japan, Kowa, Otsuka, Pfizer Japan, Rohto, Senju), Lecture fees (Alcon Japan, AMO Japan, CREWT Medical Systems, Johnson & Johnson, Kowa, MSD Japan, Novartis Pharma, Otsuka, Pfizer Japan, R-Tech Ueno, Santen, Senju).

The original article can be found online at <https://doi.org/10.1007/s10384-017-0512-2>.

✉ Kyoko Ishida
kyoko.ish@gmail.com

¹ Department of Ophthalmology, Toho University
 Ohashi Medical Center, 2-17-6 Ohashi, Meguro-ku,
 Tokyo 153-8515, Japan

² Department of Ophthalmology, Gifu University Graduate
 School of Medicine, Gifu, Japan