Killing forms on quaternion-Kähler manifolds

Andrei Moroianu · Uwe Semmelmann

Received: 9 April 2008 / Accepted: 15 April 2008 / Published online: 18 July 2008
© Springer Science+Business Media B.V. 2008

Erratum to: Ann Glob Anal Geom DOI 10.1007/s10455-005-1150-3

The aim of this note is to fill a gap in the proof of Theorem 6.1 in [1]—stating that every Killing p -form ($p \geq 2$) on a compact quaternion-Kähler manifold M^{4m} ($m \geq 2$) is parallel. This gap, pointed out by Liana David, is due to two wrong coefficients in the formulas at the middle of page 329. The correct equations read

$$\begin{aligned}pd^+u &= d(L^-u), & (p-1)d^cu &= d(Ju + 3u), & pd^-u &= d(\Lambda^+u) - \delta^cu, \\(p-1)\delta^+u &= -2d(Cu), & (p-1)\delta^cu &= -2d(\Lambda^+u) - 2d^-u, & (p-3)\delta^-u &= -2d(\Lambda u).\end{aligned}$$

The remaining part of the proof works verbatim for $p > 3$, but an extra argument is needed for $p = 2$ and $p = 3$.

The case $p = 2$. From the six equations above, one obtains the vanishing of d^+u , δ^-u , and $\delta^+u + d^cu + 3du$, but no longer that of δ^cu and d^-u . Correspondingly, the proof of Lemma 6.3 fails. Fortunately, an ad hoc argument shows that $du = 0$ in this case. Indeed, the third equation of the system (7) shows that du is an eigenform of $2C - J$ for the eigenvalue 9. On the other hand, Lemma 5.1, Lemma 5.2, and the decomposition

$$\Lambda^3(H \otimes E) = H \otimes [\Lambda_0^{2,1}E \oplus E] \oplus \text{Sym}^3 H \otimes [\Lambda_0^3E \oplus E]$$

show that the eigenvalues of $2C - J$ on the four summands of $\Lambda^3 M$ are 3, $4m + 5$, 15, and $4m + 11$, respectively. For $m \geq 2$, none of them equals 9.

The case $p = 3$. The proof works well in this case, except that one does not obtain $\delta^-u = 0$ in Lemma 6.2. However, this relation is not needed until the point (b) at the bottom of

The online version of the original article can be found under doi:[10.1007/s10455-005-1150-3](https://doi.org/10.1007/s10455-005-1150-3).

A. Moroianu (✉) · U. Semmelmann
CMLS, Ecole Polytechnique, Palaiseau 91128, France
e-mail: andrei.moroianu@math.polytechnique.fr

page 331. In order to rule out that case, one has to replace the argument given there with the fact that for $p = 3$ and $m \geq 2$, the inequality $p \geq 2m + 1$ is impossible.

Remark 1 The assumption that $m \geq 2$ is essential. For $m = 1$, the quaternionic projective space $\mathbb{H}P^1 \simeq S^4$ carries non-parallel Killing 2-forms and 3-forms (cf. [2]).

References

1. Moroianu, A., Semmelmann, U.: Twistor forms on quaternion-Kähler manifolds. *Ann. Glob. Anal. Geom.* **28**, 319–335 (2005)
2. Semmelmann, U.: Conformal Killing forms on Riemannian manifolds. *Math. Z.* **245**, 503–527 (2003)