CORRECTION

Correction to: Taxonomic resolution and functional traits in the analysis of tropical oribatid mite assemblages

Maria A. Minor¹ · Sergey G. Ermilov² · Alexei V. Tiunov^{3,4}

Published online: 4 December 2017

© Springer International Publishing AG, part of Springer Nature 2017

Correction to: Exp Appl Acarol <https://doi.org/10.1007/s10493-017-0190-2>

Due to an unfortunate turn of events, the statistical results for TS1 (F - and p -values) in Table 4 were displayed in reversed order. The correct representation of Table 4 is published here and should be treated as definitive.

The original article can be found online at <https://doi.org/10.1007/s10493-017-0190-2>.

✉ Maria A. Minor
m.a.minor@massey.ac.nz

Sergey G. Ermilov
ermilovacari@yandex.ru

Alexei V. Tiunov
a_tiunov@mail.ru

¹ Institute of Agriculture and Environment, Massey University,
Private Bag 11222, Palmerston North, New Zealand

² Tyumen State University, Tyumen, Russia

³ A.N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, Leninsky pr. 33,
Moscow, Russia 119071

⁴ The Russian-Vietnamese Tropical Scientific and Technological Centre, Southern Branch,
Ho Chi Minh, Vietnam

Table 4 PERMANOVA results for testing the hypotheses of no difference in soil Oribatida assemblages (mixed model with site as a random factor nested within forest type, $\alpha = 0.05$), Dong Nai Biosphere Reserve, Vietnam

Factor	df	Pseudo- <i>F</i>	<i>p</i>
Species			
Sampling date (wet vs. dry season)	1,12	1.32	0.20
Forest type (plantation vs. other)	1,6	2.34	0.042
Region (Dong Nai vs. Cat Tien)	1,12	2.78	0.005
Fam			
Forest type	1,6	1.89	0.063
Region	1,12	2.38	0.027
TS1			
Forest type	1,6	2.09	0.029
Region	1,12	1.55	0.18
TS2			
Forest type	1,6	2.71	0.098
Region	1,12	1.41	0.26