Correction to: The Iodide and Iodate Distribution in the Seto Inland Sea, Japan

Kazuhiko Takeda¹ · Kengo Yamane² · Yuuta Horioka² · Kazuaki Ito^{2,3}

Published online: 8 March 2018

© Springer Science+Business Media B.V., part of Springer Nature 2018

Correction to: Aquat Geochem (2017) 23:315–330

<https://doi.org/10.1007/s10498-017-9324-8>

In the original publication of the article, Figs. 3 and 4 were interchanged. Now the correct figures have been provided in this erratum (Figs. 3, 4).

The original article can be found online at <https://doi.org/10.1007/s10498-017-9324-8>.

✉ Kazuhiko Takeda
takedaq@hiroshima-u.ac.jp

Kazuaki Ito
itok@hiro.kindai.ac.jp

¹ Graduate School of Biosphere Science, Hiroshima University, 1-7-1 Kagamiyama, Higashi-Hiroshima 739-8521, Japan

² Cluster of Biotechnology and Chemistry Systems, Graduate School of Systems Engineering, Kindai University, 1 Umenobe, Takaya, Higashi-Hiroshima 739-2116, Japan

³ Department of Biotechnology and Chemistry, Faculty of Engineering, Kindai University, 1 Umenobe, Takaya, Higashi-Hiroshima 739-2116, Japan

Fig. 3 Vertical profiles of iodide, iodate, total inorganic iodine (iodide + iodate), salinity, rationalized iodide, rationalized total inorganic iodine, and the ratio of iodide to total inorganic iodine during the cruise from June 8–12, 2015. Salinities were obtained from the CTD system of the ship. The water depth was 16.0 m for OS-1, 19.7 m for OS-2, 35.9 m for OS-3, 36.0 m for OS-4, 64.5 m for KI-1, and 69.1 m for KI-2, respectively

Fig. 4 Vertical profiles of iodide, iodate, total inorganic iodine (iodide + iodate), salinity, rationalized iodide, rationalized total inorganic iodine, and the ratio of iodide to total inorganic iodine during the cruise from August 8–9, 2015. Salinities were obtained from the CTD system of the ship. The water depth was 16.5 m for HR-1, 23.7 m for HR-2, 35.3 m for HR-3, and 36.3 m for HR-4, respectively