Real-time fluorogenic PCR assays for the detection of *entA*, the gene encoding staphylococcal enterotoxin A

Jennifer R. Horsmon · Cheng J. Cao ·
Akbar S. Khan · Mark V. Gostomski ·
James J. Valdes · Kevin P. O'Connell

Published online: 10 March 2007
© Springer Science+Business Media B.V. 2007

Erratum to: Biotechnol Lett (2006) 28: 823–829
DOI 10.1007/s10529-006-9011-0

There was an error in Table 1 of the original article (the probe and primer sequences did not align to the *entA* target sequence). The corrected Table 1 is shown on the following page.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10529-006-9011-0>

J. R. Horsmon · C. J. Cao · A. S. Khan ·
J. J. Valdes · K. P. O'Connell (✉)
U.S. Army Edgewood Chemical Biological Center,
AMSRD-ECB-RT-BM, Aberdeen Proving Ground,
5183, Edgewood, MD 21010, USA
e-mail: kevin.oconnell1@us.army.mil

M. V. Gostomski
SAIC, Aberdeen Proving Ground, P.O. Box 68,
Edgewood, MD 21010, USA

C. J. Cao
US Army Center for Health Promotion and
Preventive Medicine, Blackhawk Road, E2100/1031,
Aberdeen Proving Ground, 5158, Edgewood,
MD 21010-5430, USA

A. S. Khan
Defense Threat Reduction Agency, Fort Belvoir,
VA 22060-6201, USA

Table 1 Primer and probe sequences for real-time fluorogenic PCR assays for the detection of the *entA* gene of *Staphylococcus aureus*

Assay	Start ^a	Probe and Primer sequences ^b	Amplicon length (bp)
1	297 397 320	Forward: TGCCCTAACGTGGACAACAA Reverse: CTGCTCCCTGCAATTCAGACT Probe: CACTTGTAATGGTAGCGAGAAAAGCGAAGAAATAA	101
2	714 812 741	Forward: CAATTTATGGCTAGACGGTAAAC Reverse: CTTGCTTGAAGATCCAACCTCC Probe: ACAGTACCTTTGGAAACGGTTAAAACGAATAAGAAAA	99

^a Position of first base in each oligonucleotide sequence relative to the *entA* sequence in GenBank accession L22566

^b All sequences listed in the 5' to 3' direction. The 5' base of each probe was labeled with 6-carboxy-fluorescein (FAM), and the 3' base of each probe was labeled with 6-carboxytetramethylrhodamine (TAMRA)