Production of biomass and recombinant human-like collagen in *Escherichia coli* processes with different CO₂ pulses

Wen J. Xue · Dai D. Fan · Longan Shang ·
Chen H. Zhu · Xiao X. Ma · Yuan Y. Yu

Published online: 23 January 2009
© Springer Science+Business Media B.V. 2009

Erratum to: Biotechnol Lett (2009) 31:221–226
DOI 10.1007/s10529-008-9852-9

Panels b and c in Fig. 2 of the original publication were incorrect. The correct Fig. 2 is shown on the following page of this erratum.

The online version of the original article can be found under doi:[10.1007/s10529-008-9852-9](https://doi.org/10.1007/s10529-008-9852-9).

W. J. Xue · D. D. Fan (✉) · C. H. Zhu ·
X. X. Ma · Y. Y. Yu

Department of Chemical Engineering and Shaanxi Key,
Laboratory of Degradable Biomedical Materials,
Northwest University, Xi'an 710069, China
e-mail: fandaiddi@nwu.edu.cn

D. D. Fan · L. Shang · C. H. Zhu · X. X. Ma
Shaanxi R&D Center of Biomaterials and Fermentation
Engineering, Xi'an 710069, China

L. Shang
College of Biological and Chemical Engineering,
Ningbo Institute of Technology, Zhejiang University,
Ningbo 315100, China

Fig. 2 Recombinant *E. coli* fermentation for HLC production under a 3-h CO_2 pulse of 20% concentration injected at: **a** the lag phase; **b** the post-lag phase of batch culture; **c** the pre-induction phase of fed-batch culture; **d** the post-induction phase of fed-batch culture. ■, DCW; □, Cell specific growth rate; ▲, Exit CO_2 concentration; ▼, HLC concentration