Erratum to: Role of Src in breast cancer cell migration and invasion in a breast cell/bone-derived cell microenvironment

Brant Pohorelic · R. Singh · S. Parkin ·
K. Koro · A.-D. Yang · C. Egan · A. Magliocco

Published online: 11 October 2011
© Springer Science+Business Media, LLC. 2011

Erratum to: Breast Cancer Res Treat
DOI 10.1007/s10549-011-1753-2

The original publication of the article has been published with incorrect values in table row “Y419/Total Src” pertaining to Fig. 1, part b. The correct values are given in the table presented below.

The online version of the original article can be found under doi:[10.1007/s10549-011-1753-2](https://doi.org/10.1007/s10549-011-1753-2).

B. Pohorelic · R. Singh · S. Parkin · K. Koro · A.-D. Yang ·
C. Egan (✉) · A. Magliocco
Departments of Oncology and Pathology and Laboratory,
Medicine, Faculty of Medicine, University of Calgary,
1331 29 St, NW, Calgary, AB T2N 4N2, Canada
e-mail: egan@ucalgary.ca

Fig. 1 Quantitation of total and phospho-Src in a panel of six breast cancer cell lines by Western blot. **a** Representative relative amounts of total and phospho-Src and actin in breast cancer cell lines. Blots for Src and phospho-Src were each run with their own actin control and the actin control shown is representative. **b** Values of total and phospho-Src by densitometric scan from panel A normalized to actin. Each Src blot was normalized to a different actin loading control, and normalized values can be compared with **a** across the rows but not down the columns. **c** Graphic representation of the values for total Src normalized to actin. **d** Graphic representation of the values of phospho-Src normalized to total Src