Erratum to: Comparison of relative versus absolute arm size change as criteria for quantifying breast cancer-related lymphedema: the flaws in current studies and need for universal methodology

Marek Ancukiewicz · Cynthia L. Miller · Melissa N. Skolny ·
Jean O'Toole · Laura E. Warren · Lauren S. Jammallo ·
Michelle C. Specht · Alphonse G. Taghian

Published online: 28 September 2012
© Springer Science+Business Media, LLC. 2012

**Erratum to: Breast Cancer Res Treat (2012) 135:145–152
DOI 10.1007/s10549-012-2111-8**

The original publication of the article has been published with an incorrect reference. The corrected reference [52] should read as given below.

Reference

52. Ward LC, Bunce IH, Cornish BH, Mirolo B, Thomas BJ, Jones LC (1992) Multifrequency bioelectrical impedance augments the diagnosis and management of lymphedema in post-mastectomy patients. *Eur J Clin Invest* 22:751–754

The online version of the original article can be found under doi:[10.1007/s10549-012-2111-8](https://doi.org/10.1007/s10549-012-2111-8).

M. Ancukiewicz · C. L. Miller · M. N. Skolny ·
L. E. Warren · L. S. Jammallo · A. G. Taghian (✉)
Department of Radiation Oncology, Massachusetts General
Hospital, 55 Fruit Street, Boston, MA 02114, USA
e-mail: ataghian@partners.org

J. O'Toole
Department of Physical and Occupational Therapy,
Massachusetts General Hospital, Boston, MA, USA

M. C. Specht
Division of Surgical Oncology, Massachusetts General Hospital,
Boston, MA, USA