Correction to: Association between lifestyle, menstrual/reproductive history, and histological factors and risk of breast cancer in women biopsied for benign breast disease

Rhonda Arthur¹ · Yihong Wang² · Kenny Ye¹ · Andrew G. Glass³ · Mindy Ginsberg¹ · Olivier Loudig¹ · Thomas Rohan¹

Published online: 11 April 2018
© Springer Science+Business Media, LLC, part of Springer Nature 2018

Correction to: Breast Cancer Res Treat (2017) 165:623–631
<https://doi.org/10.1007/s10549-017-4347-9>

The third category for extent of involution in Table 4 was published incorrectly in the original publication. The correct classification is $\geq 75\%$ and the corrected Table 4 is given below.

Table 4 Association between lobular involution and breast cancer risk among women who were biopsied for benign breast disease

	No. of cases	No. of controls	Crude OR (95% CI)	Multivariable OR (95% CI) ^a
Lobular involution				
No	105	85	1.00 ^b	1.00 ^b
Yes	204	215	0.90 (0.76–1.06)	1.04 (0.66–1.63)
Extent of involution				
Absent	105	85	1.00 ^b	1.00 ^b
1–74%	165	167	0.78 (0.53–1.14)	1.06 (0.67–1.69)
$\geq 75\%$	39	48	0.61 (0.35–1.07)	0.91 (0.45–1.84)
P trend			0.25	0.72

^aAdjusted for smoking history, BMI, age at menarche, age at first live birth and number of pregnancies, menopausal status, history of oophorectomy, family history, hormonal therapy use and benign breast histology (non-proliferative, hyperplasia without atypia, hyperplasia with atypia)

^bReference category

The original article can be found online at <https://doi.org/10.1007/s10549-017-4347-9>.

✉ Rhonda Arthur
rhonda.arthur@einstein.yu.edu

¹ Department of Epidemiology and Population Health, Albert Einstein College of Medicine, 1300 Morris Park Ave., Bronx, NY 10461, USA

² Department of Pathology and Laboratory Medicine, Rhode Island Hospital and Lifespan Medical Center, Warren Alpert Medical School of Brown University, Providence, RI 02903, USA

³ Center for Health Research, Kaiser Permanente Northwest, Portland, OR, USA