ERRATUM

Erratum to: Risk of breast, endometrial, colorectal, and renal cancers in postmenopausal women in association with a body shape index and other anthropometric measures

Geoffrey C. Kabat¹ · Xiaonan Xue¹ · Victor Kamensky¹ · Dorothy Lane² · Jennifer W. Bea³ · Chu Chen⁴ · Lihong Qi⁵ · Marcia L. Stefanick⁶ · Rowan T. Chlebowski⁷ · Jean Wactawski-Wende⁸ · Sylvia Wassertheil-Smoller¹ · Thomas E. Rohan¹

Published online: 27 July 2017
© Springer International Publishing AG 2017

**Erratum to: Cancer Causes Control (2015)
26:219–229
DOI 10.1007/s10552-014-0501-4**

Table 5 has been published incorrectly in the original publication of the article. The correct version of the Table 5 is provided below.

Bolded numbers in Table 5 have been corrected; the unbolded numbers were correct.

Breast cancer: age (continuous), alcohol (servings per week—continuous), smoking (pack-years—continuous), MET-hrs/wk (continuous), age at menarche (<12, 12, 13,

The online version of the original article can be found under doi:[10.1007/s10552-014-0501-4](https://doi.org/10.1007/s10552-014-0501-4).

✉ Geoffrey C. Kabat
geoffrey.kabat@einstein.yu.edu

¹ Department of Epidemiology and Population Health, Albert Einstein College of Medicine, 1300 Morris Park Avenue, Bronx, NY 10461, USA

² Department of Preventive Medicine, School of Medicine, Stony Brook University, Stony Brook, NY, USA

³ Department of Medicine, Arizona Cancer Center, Tucson, AZ, USA

⁴ Department of Epidemiology, School of Public Health, Fred Hutchinson Cancer Research Center, Seattle, WA, USA

⁵ Department of Public Health Sciences, School of Medicine, University of California at Davis, Davis, CA, USA

⁶ Department of Medicine, Stanford University, Stanford, CA, USA

⁷ Los Angeles Biomedical Research Institute at Harbor-UCLA Medical Center, Torrance, CA, USA

⁸ Department of Social and Preventive Medicine, School of Public Health and Health Professions, University at Buffalo, Buffalo, NY, USA

Type of cancer	Never hormone users (N = 62,392)	Ever hormone users (N = 81,508)
Overall	1.00	1.00
Breast	1.00	1.00
ER+	1.00	1.00
ER-	1.00	1.00
Colorectal	1.00	1.00
Prostate	1.00	1.00
Pancreatic	1.00	1.00
Stomach	1.00	1.00
Lung	1.00	1.00
Ovarian	1.00	1.00
Uterine	1.00	1.00
Cervix	1.00	1.00
Bladder	1.00	1.00
Kidney	1.00	1.00
Liver	1.00	1.00
Esophagus	1.00	1.00
Small intestine	1.00	1.00
Rectum	1.00	1.00
Vagina	1.00	1.00
Penis	1.00	1.00
Testis	1.00	1.00
Thyroid	1.00	1.00
Salivary gland	1.00	1.00
Nerve	1.00	1.00
Soft tissue	1.00	1.00
Unknown	1.00	1.00

	ABSI quintiles		BMI quintiles		WC quintiles		ABSI quintiles		BMI quintiles		WC quintiles	
	HR	95% CI	HR	95% CI	HR	95% CI	HR	95% CI	HR	95% CI	HR	95% CI
Breast (N = 7,039)	1.00	Ref.	1.00	Ref.	1.00	Ref.	1.00	Ref.	1.00	Ref.	1.00	Ref.
	1.17	1.03–1.32	1.22	1.06–1.41	1.23	1.07–1.42	0.92	0.84–1.02	1.04	0.95–1.14	1.07	0.97–1.17
	0.99	0.87–1.12	1.30	1.14–1.50	1.44	1.25–1.65	0.98	0.89–1.08	1.05	0.95–1.16	1.07	0.98–1.18
	1.04	0.92–1.19	1.50	1.31–1.72	1.52	1.33–1.57	0.99	0.90–1.09	1.11	1.01–1.23	1.14	1.04–1.26
	1.11	0.97–1.25	1.69	1.48–1.94	1.67	1.05–1.61	1.01	0.91–1.11	1.28	1.15–1.42	1.25	1.12–1.38
	P for linear trend											
Endometrium (N = 1,157)	0.45		<0.0001		<0.0001		0.50		<0.0001		<0.0001	
	1.00	Ref.	1.00	Ref.	1.00	Ref.	1.00	Ref.	1.00	Ref.	1.00	Ref.
	1.01	0.75–1.36	1.38	0.90–2.11	1.70	1.12–2.60	1.40	1.11–1.77	0.84	0.66–1.05	0.85	0.68–1.08
	1.08	0.81–1.46	2.21	1.49–3.27	1.98	1.30–3.00	1.20	0.94–1.54	0.82	0.64–1.05	0.87	0.68–1.10
	1.19	0.89–1.60	2.85	1.95–4.16	3.79	2.58–5.52	0.86	0.66–1.14	0.93	0.73–1.20	0.94	0.73–1.21
	1.42	1.07–1.88	6.15	4.30–8.80	6.03	4.16–8.76	0.98	0.74–1.29	1.16	0.89–1.51	1.14	0.88–1.48
Colorectal (N = 1,908)	P for linear trend											
	0.005		<0.0001		<0.0001		0.14		0.19		0.30	
	1.00	Ref.	1.00	Ref.	1.00	Ref.	1.00	Ref.	1.00	Ref.	1.00	Ref.
	0.99	0.79–1.26	0.95	0.77–1.19	1.40	1.10–1.78	1.19	0.95–1.50	1.45	1.16–1.80	1.57	1.25–1.97
	1.26	1.01–1.57	0.88	0.71–1.10	1.21	0.95–1.54	1.35	1.08–1.69	1.48	1.18–1.85	1.53	1.22–1.94
	1.22	0.98–1.52	1.05	0.85–1.30	1.45	1.15–1.83	1.29	1.03–1.62	1.52	1.21–1.90	1.93	1.54–2.43
	1.47	1.19–1.82	1.25	1.02–1.54	1.81	1.44–2.28	1.49	1.19–1.87	1.60	1.26–2.03	1.99	1.56–2.53
	P for linear trend											
<0.0001		0.003		<0.0001		0.0006		0.0009		<0.0001		

≥ 14), age at first birth (<20 , $20-29$, ≥ 30 , missing), parity (continuous), family history of breast cancer in a first degree relative (yes, no, missing), history of breast biopsy (yes, no, missing), education (less than high school grad, high school grad/some college, college grad, post-college), ethnicity (white, black, other), treatment allocation (observational study; treatment/control/placebo arm of clinical trials).

Endometrial cancer: age, alcohol, smoking, parity, oral contraceptive use (ever, never), education, ethnicity, treatment allocation.

Colorectal cancer: age (continuous), alcohol, smoking, MET-hrs/wk (continuous), aspirin intake, diabetes, family history of colorectal cancer in a first degree relative, education, ethnicity, treatment allocation.

Last paragraph of Results section should read:

Among never-hormone users, after mutual adjustment using residuals, the HRs for breast cancer and endometrial cancer for the highest relative to the lowest quintile of BMI and WC were weakened compared to the values reported in Table 5 (i.e., compared to the new, corrected values): for breast cancer, $HR_{BMI} = 1.39$, 95% CI 1.13–1.72; $HR_{WC} = 1.30$, 95% CI 1.05–1.61, respectively; for endometrial cancer: $HR_{BMI} = 2.96$, 95% CI 1.71–5.12; $HR_{WC} = 2.53$, 95% CI 1.44–4.44. After mutual adjustment, BMI was inversely associated with colorectal cancer in never hormone users: $HR_{BMI} = 0.62$, 95% CI 0.44–0.89, whereas the positive association with WC remained: $HR_{WC} = 2.19$, 95% CI 1.54–3.13.