It Is Possible to Classify Non-erosive Reflux Disease (NERD) Patients into Endoscopically Normal Groups and Minimal Change Groups by Subjective Symptoms and Responsiveness to Rabeprazole—A Report from a Study with Japanese Patients

Motoyasu Kusano · Naohito Shirai · Kanako Yamaguchi ·
Yasuyuki Shimoyama · Michio Hongo · Tsutomu Chiba ·
Yoshikazu Kinoshita · The Acid-Related Symptom (ARS) Research Group

Published online: 2 December 2008
© Springer Science+Business Media, LLC 2008

Erratum to: Dig Dis Sci
DOI 10.1007/s10620-008-0290-2

This article was published with an incomplete author listing—Yasuyuki Shimoyama was left out. This erratum includes the name and affiliation of the missing author.

The online version of the original article can be found under doi:[10.1007/s10620-008-0290-2](https://doi.org/10.1007/s10620-008-0290-2).

M. Kusano (✉) · Y. Shimoyama
Department of Endoscopy and Endoscopic Surgery, Gunma
University Hospital, 3-39-15 Showamachi, Maebashi, Gunma
371-8511, Japan
e-mail: mkusano@showa.gunma-u.ac.jp

N. Shirai
Department of Gastroenterology, Enshu Hospital,
Hamamatsu, Japan

K. Yamaguchi
Department of Internal Medicine, Saga Medical School,
Saga, Japan

M. Hongo
Departments of Comprehensive Medicine and Psychosomatic
Medicine, Tohoku University Hospital, Sendai, Japan

T. Chiba
Department of Gastroenterology and Hepatology, Kyoto
University Graduate School of Medicine, Kyoto, Japan

Y. Kinoshita
Department of Gastroenterology and Hepatology, Shimane
University School of Medicine, Izumo, Japan

The Acid-Related Symptom (ARS) Research Group
Izumo, Japan