Is optic nerve fibre mis-routing a feature of congenital stationary night blindness?

**T. Ung · L. E. Allen · A. T. Moore · D. Trump ·
I. Zito · A. J. Hardcastle · J. Yates ·
K. Bradshaw**

Published online: 13 July 2006
© Springer Science+Business Media B.V. 2006

In Documenta Ophthalmologica (2005) 111: 169–178 the article ‘*Is optic nerve fiber mis-routing a feature of congenital stationary night blindness?*’ has been published. Unfortunately, the list of authors was incomplete. Please find the complete list including affiliations here:

T. Ung¹, L.E. Allen¹, A.T. Moore^{1,2}, D. Trump^{3,4}, I. Zito⁵, A.J. Hardcastle⁵, J. Yates³, & K. Bradshaw¹

¹Ophthalmology Department, Addenbrooke’s Hospital, Cambridge, CB2 2QQ, UK;

²Moorfields Eye Hospital, London, UK;

³Genetics Department, Addenbrooke’s Hospital, Cambridge, UK;

⁴Clinical Genetics, University of Manchester, Manchester, UK;

⁵Division of Molecular and Cellular Neuroscience, Institute of Ophthalmology, University College, London, UK

The online version of the original article can be found
at <http://dx.doi.org/10.1007/s10633-005-5503-9>

T. Ung · L. E. Allen · A. T. Moore · K. Bradshaw (✉)
Ophthalmology Department, Addenbrooke’s
Hospital, Cambridge CB2 2QQ, UK
e-mail: keith.bradshaw@addenbrookes.nhs.uk

A. T. Moore
Moorfields Eye Hospital, London, UK

D. Trump · J. Yates
Genetics Department, Addenbrooke’s Hospital,
Cambridge, UK

D. Trump
Clinical Genetics, University of Manchester,
Manchester, UK

I. Zito · A. J. Hardcastle
Division of Molecular and Cellular Neuroscience,
Institute of Ophthalmology, University College,
London, UK