Evaluation of pastures for horses grazing on soils polluted by trace elements

P. Madejón · M. T. Domínguez · J. M. Murillo

Published online: 10 March 2009
© Springer Science+Business Media, LLC 2009

Erratum to: Ecotoxicology DOI 10.1007/s10646-009-0296-3

The author regrets for error in Table 2 of his article. Correct version of Table 2 is as given below.

Table 2 Species composition of the pastures (affected and control sites)

Family	Sp.	Spring	Autumn	Family	Sp.	Spring	Autumn
Poaceae	<i>Agrostis pourretii</i>	0.08	0.35	Fabaceae	<i>Biserrula pelecinus</i>	0.01	
	<i>Avena barbata</i> subsp. <i>barbata</i>	0.03			<i>Medicago doliata</i>	0.06	0.01
	<i>Avena sterilis</i>	0.1	0.05		<i>Medicago murex</i>	0.07	0.01
	<i>Bromus diandrus</i>	0.17	0.11		<i>Medicago polymorpha</i>	0.13	0.03
	<i>Bromus lanceolatus</i>	0.21			<i>Scorpiurus muricatus</i>	0.04	0.04
	<i>Bromus madritensis</i>	0.1			<i>Vicia villosa</i> subsp. <i>varia</i>	0.02	
	<i>Cynodon dactylon</i>	0.01	0.25	Others	<i>Anagallis arvensis</i>	0.07	0.01
	<i>Hordeum marinum</i>	0.06			<i>Beta vulgaris</i>	0.01	
	<i>Lamarckia aurea</i>	0.03			<i>Convulvulus arvensis</i>	0.01	0.06
	<i>Lolium perenne</i>	0.1			<i>Echium plantagineum</i>		0.05
Asteraceae	<i>Phalaris minor</i>	0.05	0.03		<i>Equisetum arvense</i>	0.01	0.03
	<i>Piptatherum miliaceum</i>	0.08	0.1		<i>Euphorbia helioscopia</i>	0.01	
	<i>Andryala integrifolia</i>	0.1			<i>Heliotropium europaeum</i>		
	<i>Carduus pycnocephalus</i>	0.02			<i>Hirschfeldia incana</i>		
	<i>Chrysanthemum coronarium</i>	0.09			<i>Malva hispanica</i>	0.04	0.05
	<i>Coleostephus myconis</i>	0.21	0.12		<i>Plantago coronopus</i>	0.01	0
	<i>Conyza bonariensis</i>	0.05			<i>Plantago lagopus</i>	0.05	0
	<i>Galactites tomentosa</i>	0.02			<i>Ranunculus trilobus</i>	0.03	0
	<i>Lactuca serriola</i>	0.01			<i>Scirpoides holoschoenus</i>		0.02
	<i>Leontodon longirostris</i>	0.09	0.02		<i>Torilis arvensis</i>	0.04	0.07
	<i>Senecio lividus</i>	0.22	0.01				
	<i>Senecio vulgaris</i>	0.14	0.16				
	<i>Silybum marianum</i>	0.03	0.02				
	<i>Xanthium strumarium</i>		0.01				

The relative frequency of observations of each species ($N = 100$) during the spring and autumn sampling is indicated

The online version of the original article can be found under doi:[10.1007/s10646-009-0296-3](https://doi.org/10.1007/s10646-009-0296-3).

P. Madejón (✉) · M. T. Domínguez · J. M. Murillo
Instituto de Recursos Naturales y Agrobiología de Sevilla (IRNAS), CSIC, P.O. Box 1052, 41080 Sevilla, Spain
e-mail: pmadejon@irnase.csic.es