

## Erratum to: Preliminary assessment of surface soil lead concentrations in Melbourne, Australia

Mark A. S. Laidlaw  · Callum Gordon · Andrew S. Ball

Published online: 11 September 2017  
© Springer Science+Business Media B.V. 2017

**Erratum to: Environ Geochem Health**  
**DOI 10.1007/s10653-017-0010-y**

Unfortunately, in the original publication of the article, Fig. 1 was incorrectly published. The original article was corrected (Fig. 1).

---


The online version of the original article can be found under doi:[10.1007/s10653-017-0010-y](https://doi.org/10.1007/s10653-017-0010-y).

---

M. A. S. Laidlaw (✉) · C. Gordon · A. S. Ball  
Centre for Environmental Sustainability and Remediation  
(EnSuRe), School of Science, RMIT University,  
PO Box 71, Bundoora, VIC 3083, Australia  
e-mail: mark.laidlaw@rmit.edu.au

C. Gordon  
e-mail: callumgordon1990@gmail.com

A. S. Ball  
e-mail: andy.ball@rmit.edu.au



**Fig. 1** Lead concentration (mg/kg) of surface soil (0–2 cm) samples collected along three transects oriented across the Melbourne metropolitan area (<250  $\mu\text{m}$  grain size). *Note* Australian NEPC Residential HIL-A Soil Pb Guideline = 300 mg/kg (NEPC 2013a, b); Australian NEPC

Residential HIL-B Soil Pb Guideline = 1200 mg/kg; and Australian NEPC Residential HIL-C Soil Pb Guideline = 600 mg/kg; California Soil Pb Guideline = 80 mg/kg (CEPA 2009)