CORRECTION

Correction to: Resistance evaluation of differentials and commercial wheat cultivars to stripe rust (*Puccinia striiformis*) infection in hot spot regions of Canada

Gurcharn Singh Brar · Raman Dhariwal ·
Harpinder Singh Randhawa

Published online: 23 March 2018
© Koninklijke Nederlandse Planteziektenkundige Vereniging 2018

Correction to: European Journal of Plant Pathology
<https://doi.org/10.1007/s10658-018-1446-3>

Table 2 AAC Penhold is CPSR wheat and carries Yr18 (not CWRS).

Fig. 2 Two-dimensional hierarchical clustering of site-years and wheat cultivars based on percent severity of stripe rust from natural infection. Colors for each cell represent infection response of a cultivar at given site-year. Here red = resistant (R), orange = moderately resistant (MR), light blue = moderately resistant to moderately susceptible (MRMS), and dark blue = susceptible (S) infection responses and Cre- Creston, British Columbia; Leth- Lethbridge, Alberta.

The online version of the original article can be found at
<https://doi.org/10.1007/s10658-018-1446-3>

G. S. Brar
Crop Development Centre/Department of Plant Science,
University of Saskatchewan, 51 Campus Dr, Saskatoon, SK S7N
5A8, Canada

R. Dhariwal · H. S. Randhawa (✉)
Lethbridge Research and Development Centre, Agriculture and
Agri-Food Canada, 5403-1 Avenue South, Lethbridge, AB T1J
4B1, Canada
e-mail: harpinder.randhawa@agr.gc.ca