Erratum to: A Banach space-valued ergodic theorem and the uniform approximation of the integrated density of states

Daniel Lenz · Fabian Schwarzenberger · Ivan Veselić

Published online: 20 September 2011
© Springer Science+Business Media B.V. 2011

Erratum to: Geom Dedicata (2011) 150:1–34 DOI 10.1007/s10711-010-9491-x

M. Keller and F. Pogorzelski pointed out that equation (10) and consequently the claim of the Ergodic Theorem 3.1 in [1] are not correct as stated. The theorem is correct, however, if a mild additional symmetry condition on the underlying group is satisfied, as we explain in what follows:

We say that $Q \subseteq G$ tiles the group G *symmetrically* if there exists a set $K \subseteq G$ satisfying $K = K^{-1}$ such that $\{Qg \mid g \in K\}$ is a disjoint partition of G .

Assumption A. G is an amenable group generated by a finite and symmetric set S , $\mathcal{A}$ is a finite set and $\mathcal{C} : G \rightarrow \mathcal{A}$ is a map, which we will call a colouring. There exists a Følner sequence $(Q_n)_{n \in \mathbb{N}}$ in G such that each element of this sequence tiles the group symmetrically. $(U_j)_{j \in \mathbb{N}}$ is a Følner sequence along which the frequencies $\nu_P = \lim_{j \rightarrow \infty} |U_j|^{-1} \sharp_P(\mathcal{C}|_{U_j})$ exist for all patterns $P \in \bigcup_{n \in \mathbb{N}} \mathcal{P}(Q_n)$. $(X, \|\cdot\|)$ is a Banach space.

We refer the reader to [1] for a detailed explanation of the notation and concepts. A complete version of the paper, where the correction of the present Erratum has been incorporated, is available as the 2nd version of the arxiv.org preprint 1003.3620.

The online version of the original article can be found under doi:[10.1007/s10711-010-9491-x](https://doi.org/10.1007/s10711-010-9491-x).

D. Lenz

Mathematisches Institut, Friedrich-Schiller-Universität Jena, 07743 Jena, Germany
URL: <http://www.analysis-lenz.uni-jena.de/>

F. Schwarzenberger (✉) · I. Veselić

Emmy-Noether-Projekt, Fakultät für Mathematik, TU Chemnitz, 09107 Chemnitz, Germany
e-mail: fabian.schwarzenberger@mathematik.tu-chemnitz.de
URL: <http://www-user.tu-chemnitz.de/~fabis>

I. Veselić

URL: <http://www.tu-chemnitz.de/mathematik/stochastik>

Remark B. The condition that G contains a Følner sequence (Q_n) such that each Q_n symmetrically tiles G is particularly satisfied if there exists a sequence of subgroups $(G_n)_{n \in \mathbb{N}}$ such that one can choose the associated fundamental domains $(Q_n)_{n \in \mathbb{N}}$ to be a Følner sequence. This is the case for any residually finite, amenable group, in particular, for any group of polynomial volume growth.

Theorem C. Assume A. For a given $\mathcal{C}$ -invariant and almost-additive function $F : \mathcal{F}(G) \rightarrow X$ the following limits

$$\bar{F} := \lim_{j \rightarrow \infty} \frac{F(U_j)}{|U_j|} = \lim_{n \rightarrow \infty} \sum_{P \in \mathcal{P}(Q_n)} \nu_P \frac{\tilde{F}(P)}{|Q_n|}$$

exist and are equal. Furthermore, for $j, n \in \mathbb{N}$ the difference

$$\Delta(j, n) := \left\| \frac{F(U_j)}{|U_j|} - \sum_{P \in \mathcal{P}(Q_n)} \nu_P \frac{\tilde{F}(P)}{|Q_n|} \right\|$$

satisfies the estimate

$$\Delta(j, n) \leq \frac{b(Q_n)}{|Q_n|} + (C + D) \frac{|\partial^{\text{diam}(Q_n)} U_j|}{|U_j|} + C \sum_{P \in \mathcal{P}(Q_n)} \left| \frac{\sharp_P(\mathcal{C}|_{U_j})}{|U_j|} - \nu_P \right|. \quad (\text{a})$$

We explain how to replace now the incorrect part of the proof of the Ergodic Theorem 3.1 in [1]. By assumption, for each fixed $n \in \mathbb{N}$, the set Q_n symmetrically tiles the group G , i.e., there exists a symmetric set $G_n \subseteq G$ such that $G = \bigcup_{g \in G_n} Q_n g$, where $Q_n g \cap Q_n h = \emptyset$ for all $g, h \in G_n$ with $g \neq h$. This property remains valid after shifting the grid by an arbitrary $x^{-1} \in G$. In fact we have

$$G = Gx^{-1} = \bigcup_{g \in G_n} Q_n g x^{-1} = \bigcup_{g \in G_n x^{-1}} Q_n g,$$

where $G_n x^{-1} := \{gx^{-1} | g \in G_n\}$. Still $Q_n g \cap Q_n h = \emptyset$ holds for all distinct $g, h \in G_n x^{-1}$, since $g = \tilde{g}x^{-1}$ and $h = \tilde{h}x^{-1}$ for some distinct $\tilde{g}, \tilde{h} \in G_n$ and

$$Q_n g \cap Q_n h = \emptyset \Leftrightarrow Q_n \tilde{g} x^{-1} \cap Q_n \tilde{h} x^{-1} = \emptyset \Leftrightarrow Q_n \tilde{g} \cap Q_n \tilde{h} = \emptyset.$$

Furthermore, we have the equality

$$\{z \in G | Q_n z \subseteq A\} = \bigcup_{x \in Q_n} \{z \in G_n x^{-1} | Q_n z \subseteq A\} \quad (\text{b})$$

since for each $z \in G$ there is $x \in Q_n$ and $g \in G_n$ with $z^{-1} = xg$ and hence $z = g^{-1}x^{-1} \in G_n x^{-1}$, as G_n is a symmetric subset of G . To see that the union in (b) is disjoint observe that for given $x, y \in Q_n$ with $x \neq y$ and $z \in G_n x^{-1}$, we have $z^{-1} \in xG_n$. By the tiling property of Q_n , this gives $z^{-1} \notin yG_n$ and hence $z \notin G_n y^{-1}$.

The $\mathcal{C}$ -invariance of F and the equation (b) imply

$$\sum_{P \in \mathcal{P}(Q_n)} \sharp_P(\mathcal{C}|_{U_j}) \tilde{F}(P) = \sum_{z \in G: Q_n z \subseteq U_j} F(Q_n z) = \sum_{x \in Q_n} \sum_{g \in I(U_j, x, n)} F(Q_n g), \quad (\text{c})$$

which replaces the incorrect formula (10) in the original paper [1].

Acknowledgments We thank M. Keller and F. Pogorzelski who pointed out an error in [1].

Reference

1. Lenz, D., Schwarzenberger, F., Veselić, I.: A Banach space-valued ergodic theorem and the uniform approximation of the integrated density of states. *Geom. Dedicata* **150**(1), 1–34 (2011)