Erratum

Growth responses of tropical cladocerans to seston from Lake Monte Alegre (Brazil) supplemented with phosphorus, fatty acids, a green algae and a cyanobacterium[☆]

A. da Silva Ferrão-Filho & M.S. Arcifa

Hydrobiologia (2006) DOI 10.1007/s10750-005-0001-3

The following corrections must be made in this article:

Page 2, right-hand column, paragraph 2, line 2: the reference should be Ferrão-Filho et al. (2003a).

Page 5: The correct Table 3 is given below.

Page 7: The correct Table 4 is given below.

Page 8, left-hand column, paragraph 1, last line: replace 'Table 1' by 'Table 2'.

The publishers apologise for these errors.

Table 3. Results of the general ANOVA for the growth bioassay in summer

Species	Treatments				df	F	p
	Seston	+PO ₄	+PUFA	+Sce			
<i>C. richardi</i>	0.183 ± 0.018(a)	0.185 ± 0.006(a)	0.146 ± 0.017(a)	0.269 ± 0.011(b)	3.8	14.3	0.0014
<i>D. gessneri</i>	0.293 ± 0.017(a)	0.255 ± 0.009(a)	0.342 ± 0.010(ab)	0.377 ± 0.003(b)	3.8	23.6	0.0003

Data are mean growth rates (d⁻¹) ± SE for both species during the four days of experiment. There were three replicate bottles per treatment, which included: (1) seston; (2) seston + PO₄; (3) seston + PUFA and (4) seston + *Scenedesmus*. In the addition treatments, 3.0 μmol P l⁻¹, 0.3 mg C l⁻¹ of PUFA-rich fish oil emulsions, and 0.5 mg C l⁻¹ of green alga were added to seston. df are the degrees of freedom for the treatments and error. Treatments with same letters are not significantly different (*post hoc* Tukey test, *p* < 0.05).

Table 4. Results of the general ANOVA for the growth bioassay in autumn

Species	Treatments				df	F	p
	Seston	+PO ₄	+Syn	+Sce			
<i>C. richardi</i>	0.313 ± 0.031(a)	0.421 ± 0.030(ab)	0.520 ± 0.027(b)	0.683 ± 0.040(c)	3.8	23.1	0.0003
<i>D. ambigua</i>	0.085 ± 0.024(a)	0.163 ± 0.018(a)	0.326 ± 0.008(b)	†	2.6	46.3	0.0002
<i>D. gessneri</i>	0.084 ± 0.037(a)	0.288 ± 0.012(b)	0.413 ± 0.019(c)	0.461 ± 0.007(c)	3.8	59.1	<0.0001
<i>M. micrura</i>	0.334 ± 0.035(a)	0.407 ± 0.018(a)	0.932 ± 0.004(b)	1.017 ± 0.028(b)	3.8	214.3	<0.0001

Data are mean growth rates (d⁻¹) ± SE during the first two days of the experiment for *C. richardi* and *M. micrura*, and during four days for *D. ambigua* and *D. gessneri*. There were three replicate bottles per treatment, which included: (1) seston; (2) seston + PO₄; (3) seston + *Synechococcus* and (4) seston + *Scenedesmus*. In the addition treatments, 3.0 μmol P l⁻¹, 0.5 mg C l⁻¹ of cyanobacterium, and 0.5 mg C l⁻¹ of green alga were added to seston. df are the degrees of freedom for the treatments and error. Treatments with same letters are not significantly different (Tukey test, *p* < 0.05). †Not performed due to lack of animals.

[☆] The online version of the original article can be found at <http://dx.doi.org/10.1007/s10750-005-0001-3>