Erratum to: A ^{57}Fe Mössbauer study on the FeSe and Fe(Se,Te) superconductors: Discontinuities in the hyperfine parameters at T_c [Hyp. Int. 208, 133 (2012).]

J. Lindén¹ · E. Rautama² · M. Karppinen² · S. Kamali³

Published online: 25 February 2016
© Springer International Publishing Switzerland 2016

**Erratum to: Hyperfine Interact (2012) 208:133136
DOI 10.1007/s10751-011-0482-6.**

Some of us reported a drop at T_c in the second-order Doppler shift and absorption in ^{57}Fe Mössbauer spectra of samples of FeSe and Fe(Se,Te). In a recent independent remeasurement of the same samples using ^{57}Fe Mössbauer spectroscopy the reported drops were no longer observed. Temporality malfunctioning at liquid-He temperatures of the cooling equipment is speculated to have caused the drop in the original data.

PACS 74.70.Xa · 76.80.+y

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10751-011-0482-6>.

✉ J. Lindén
jlinden@abo.fi

¹ Physics Department, Åbo Akademi University, FIN-20550 Turku, Finland

² Department of Chemistry, Aalto University School of Science and Technology, FI-00076 Aalto, Finland

³ Department of Mechanical, Aerospace, and Biomechanical Engineering, University of Tennessee Space Institute, Tullahoma, TN 37388, USA