



## Correction to: Characterization of *Mycobacterium tuberculosis* ferredoxin with Mössbauer spectroscopy

Christina S. Müller<sup>1</sup> · Dominique F. Bechtel<sup>2</sup> · Hendrik Auerbach<sup>1</sup> · Juliusz A. Wolny<sup>1</sup> · Antonio J. Pierik<sup>2</sup> · Volker Schünemann<sup>1</sup>

Published online: 10 February 2020  
© Springer Nature Switzerland AG 2020

**Correction to: *Hyperfine Interactions* (2019) 240: 117**  
<https://doi.org/10.1007/s10751-019-1678-4>

The article was published with erroneous values in Table 1. Please find in this document the correct version of Table 1 that should be regarded as the final version by the reader. The following changes have been made to Table 1: The numerical values of all A-Tensors are given in kG (kilo Gauss) instead of T (Tesla). Therefore the unit is changed from T to kG in the corrected version.

---

This article is part of the Topical Collection on *Proceedings of the International Conference of the Application of the Mössbauer Effect (ICAME 2019) held in Dalian, China, 1-6 September 2019*  
Edited by Tao Zhang, Junhu Wang and Xiaodong Wang

---

The online version of the original article can be found at <https://doi.org/10.1007/s10751-019-1678-4>

✉ Christina S. Müller  
csmueller@physik.uni-kl.de

Antonio J. Pierik  
pierik@chemie.uni-kl.de

<sup>1</sup> Department of Physics, Technische Universität Kaiserslautern, Erwin-Schrödinger-Str. 46, 67663 Kaiserslautern, Germany

<sup>2</sup> Department of Chemistry, Technische Universität Kaiserslautern, Erwin-Schrödinger-Str. 54, 67663 Kaiserslautern, Germany

**Table 1** Mössbauer parameters of *Mt* Fdx at  $T = 4.2$  K compared to the reduced form of *Dg* Fdx II at  $T = 4.2$  K [10, 15]

Component	<i>Mt</i> Fdx		<i>Dg</i> Fdx II [9,15]	
	1	2	1	2
$\delta$ (mms <sup>-1</sup> )	0.27 ( $\pm 0.02$ )	0.44 ( $\pm 0.02$ )	0.32	0.46
$\Delta E_Q$ (mms <sup>-1</sup> )	-0.53 ( $\pm 0.03$ )	1.47 ( $\pm 0.03$ )	-0.52	1.47
$\Gamma$ (mms <sup>-1</sup> )	0.33* ( $\pm 0.02$ )	0.33* ( $\pm 0.02$ )	-	-
$\eta$	-2 ( $\pm 0.2$ )	0.4 ( $\pm 0.2$ )	-2	0.4
$D$ (cm <sup>-1</sup> )	-1.70 ( $\pm 0.1$ )	-1.70 ( $\pm 0.1$ )	-2.5	-2.5
$E/D$	0.33 (-0.05)	0.33 (-0.05)	0.23	0.23
$A/\mu_{\text{NGN}}$ (kG)	(116/120/129) ( $\pm 5$ )	(-140/-140/-126) ( $\pm 5$ )	(99/116/126)	(-149/-149/-116)
% Area	33	67	33	67

\*) For the simulation of the spectrum at  $T = 77$   $\Gamma_1 = 0.30$  mms<sup>-1</sup> and  $\Gamma_2 = 0.35$  mms<sup>-1</sup> was used

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.