Impact of Gun-hunting on Diurnal Primates in Continental Equatorial Guinea

Noëlle F. Kümpel · E. J. Milner-Gulland ·
Guy Cowlshaw · J. Marcus Rowcliffe

Published online: 9 July 2008
© Springer Science + Business Media, LLC 2008

Due to a publication error, a corrupted version of Fig. 1 was published in the article. We regret the error and publish a corrected version of the figure here.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10764-008-9254-9>.

N. F. Kümpel (✉)
Zoological Society of London, Regent's Park, London NW1 4RY, UK
e-mail: noelle.kumpel@zsl.org

N. F. Kümpel
Department of Anthropology, University College London, 14 Taviton Street,
London WC1H 0BW, UK

N. F. Kümpel · E. J. Milner-Gulland
Division of Biology, Imperial College London, Silwood Park Campus, Ascot,
Berkshire SL5 7PY, UK

G. Cowlshaw · J. M. Rowcliffe
Institute of Zoology, Zoological Society of London, Regent's Park, London NW1 4RY, UK

Fig. 1 Study area, showing hunter camps, hunter follows, and location of transects. Sendje is demarcated by a black triangle, hunter camps by large black dots, subject encounters during hunter follows by small gray dots, and transects by black lines. Transect zones (calculated as the area of a minimum convex polygon encircling the group of transects in each site, including a 500-m buffer to account for typical *Colobus satanus* home range size of ≥ 30 ha: Gautier-Hion *et al.* 1999) are circled. The darker gray area is Monte Alén National Park.