0-Hecke algebra actions on coinvariants and flags

Jia Huang

Received: 13 December 2012 / Accepted: 28 October 2013 / Published online: 19 November 2013
© Springer Science+Business Media New York 2013

Abstract The 0-Hecke algebra $H_n(0)$ is a deformation of the group algebra of the symmetric group $\mathfrak{S}_n$. We show that its coinvariant algebra naturally carries the regular representation of $H_n(0)$, giving an analogue of the well-known result for $\mathfrak{S}_n$ by Chevalley–Shephard–Todd. By investigating the action of $H_n(0)$ on coinvariants and flag varieties, we interpret the generating functions counting the permutations with fixed inverse descent set by their inversion number and major index. We also study the action of $H_n(0)$ on the cohomology rings of the Springer fibers, and similarly interpret the (non-commutative) Hall–Littlewood symmetric functions indexed by hook shapes.

Keywords 0-Hecke algebra · Coinvariant algebra · Demazure operator · Descent monomial · Flag variety · Hall–Littlewood function · Ribbon number · Springer fiber

1 Introduction

A composition α of n gives rise to a descent class of permutations in $\mathfrak{S}_n$; the cardinality of this descent class is known as the *ribbon number* r_α and its inv-generating function is the *q -ribbon number* $r_\alpha(q)$. Reiner and Stanton [27] defined a *(q, t) -ribbon number* $r_\alpha(q, t)$, and gave an interpretation by representations of the symmetric group $\mathfrak{S}_n$ and the finite general linear group $GL(n, \mathbb{F}_q)$. The main goal of this work is to obtain similar interpretations of various ribbon numbers by representations of the 0-Hecke algebra $H_n(0)$ of type A , a deformation of the group algebra of $\mathfrak{S}_n$ over an arbitrary field $\mathbb{F}$. It is defined as an associative $\mathbb{F}$ -algebra generated by

J. Huang (✉)

School of Mathematics, University of Minnesota, Minneapolis, MN 55455, USA
e-mail: huang338@umn.edu

$T_1, \dots, T_{n-1}$ with relations

$$\begin{cases} T_i^2 = -T_i, & 1 \leq i \leq n-1, \\ T_i T_{i+1} T_i = T_{i+1} T_i T_{i+1}, & 1 \leq i \leq n-2, \\ T_i T_j = T_j T_i, & |i-j| > 1. \end{cases}$$

It has an $\mathbb{F}$ -basis $\{T_w : w \in \mathfrak{S}_n\}$ where $T_w := T_{i_1} \cdots T_{i_k}$ for any reduced expression $w = s_{i_1} \cdots s_{i_k}$.

When $\mathbb{F}$ is an algebraically closed field of characteristic zero, it is well-known that the simple $\mathfrak{S}_n$ -modules S_λ are indexed by partitions $\lambda \vdash n$, and every $\mathfrak{S}_n$ -module is a direct sum of simple $\mathfrak{S}_n$ -modules; the *Frobenius characteristic map* ch sends a direct sum of simple $\mathfrak{S}_n$ -modules to the sum of the corresponding Schur functions s_λ . There is an analogue for $H_n(0)$ -modules which holds over an arbitrary field $\mathbb{F}$. Norton [26] showed that

$$H_n(0) = \bigoplus_{\alpha \models n} \mathbf{P}_\alpha \quad (1)$$

summed over all compositions α of n , where every $\mathbf{P}_\alpha$ is a (left) indecomposable $H_n(0)$ -module. It follows that $\{\mathbf{P}_\alpha : \alpha \models n\}$ is a complete list of non-isomorphic projective indecomposable $H_n(0)$ -modules, and $\{\mathbf{C}_\alpha : \alpha \models n\}$ is a complete list of non-isomorphic simple $H_n(0)$ -modules, where $\mathbf{C}_\alpha = \text{top}(\mathbf{P}_\alpha) = \mathbf{P}_\alpha / \text{rad } \mathbf{P}_\alpha$ for all compositions $\alpha \models n$.

By Krob and Thibon [21], the finitely generated $H_n(0)$ -modules correspond to the quasi-symmetric functions via the *quasi-symmetric characteristic* Ch which sends a simple $\mathbf{C}_\alpha$ to the *fundamental quasi-symmetric function* F_α , and the finitely generated projective $H_n(0)$ -modules correspond to the non-commutative symmetric functions via the *non-commutative characteristic* $\mathbf{ch}$ which sends a projective indecomposable $\mathbf{P}_\alpha$ to the *non-commutative ribbon Schur function* $\mathbf{s}_\alpha$. There are also graded versions of the two characteristic maps Ch and $\mathbf{ch}$ for finitely generated (projective) $H_n(0)$ -modules with filtrations.

The symmetric group $\mathfrak{S}_n$ acts on the polynomial ring $\mathbb{F}[X] := \mathbb{F}[x_1, \dots, x_n]$ by permuting the variables, and the 0-Hecke algebra $H_n(0)$ acts on $\mathbb{F}[X]$ via the *Demazure operators* $\pi_1, \dots, \pi_{n-1}$, where

$$\pi_i f := \frac{x_i f - x_{i+1} s_i(f)}{x_i - x_{i+1}}, \quad \forall f \in \mathbb{F}[X]. \quad (2)$$

The coinvariant algebra of $H_n(0)$ coincides with that of $\mathfrak{S}_n$, since $\pi_i f = f$ if and only if $s_i f = f$ for all $f \in \mathbb{F}[X]$. The following result for $\mathfrak{S}_n$ is well known.

Theorem 1.1 (Chevalley–Shephard–Todd) *The coinvariant algebra $\mathbb{F}[X]/(\mathbb{F}[X]_+^{\mathfrak{S}_n})$ is isomorphic to the regular representation of $\mathfrak{S}_n$, if $\mathbb{F}$ is a field of characteristic 0.*

We give an analogue for the 0-Hecke algebras.

Theorem 1.2 *The coinvariant algebra $\mathbb{F}[X]/(\mathbb{F}[X]_+^{\mathfrak{S}_n})$ with $H_n(0)$ -action defined by (2) is isomorphic to the regular representation of $H_n(0)$, where $\mathbb{F}$ is an arbitrary field.*

We prove this theorem by showing a similar decomposition to Norton's decomposition (1). This leads to an $\mathbb{F}$ -basis of the coinvariant algebra, which is closely related to the well-known basis of *descent monomials*. Our new basis consists of certain *Demazure atoms* obtained by consecutively applying the operators $\bar{\pi}_i = \pi_i - 1$ to some descent monomials. Theorem 1.2 and its proof are also valid when $\mathbb{F}$ is replaced with $\mathbb{Z}$.

It follows from Theorem 1.2 that the coinvariant algebra has not only the grading by the degrees of polynomials, but also the filtration by the length of permutations in $\mathfrak{S}_n$. This completes the following picture.

Here the inverse descent set $D(w^{-1})$ is identified with the composition α of n with $D(\alpha) = D(w^{-1})$. We shall see in Sect. 3 that r_α and $r_\alpha(q)$ appear as coefficients of F_α in $\text{Ch}(H_n(0))$ and $\text{Ch}_q(H_n(0))$, respectively.

Next we consider the finite general linear group $G = GL(n, \mathbb{F}_q)$, where q is a power of a prime p , and its Borel subgroup B . The 0-Hecke algebra $H_n(0)$ acts on the flag variety $1_B^G = \mathbb{F}[G/B]$ by $T_w B = BwB$ if $\text{char}(\mathbb{F}) = p$, and this induces an action on the coinvariant algebra $\mathbb{F}[X]^B / (\mathbb{F}[X]_+^G)$ of the pair (G, B) (see Sect. 7). By studying the (graded) multiplicities of the simple $H_n(0)$ -modules in these $H_n(0)$ -modules, we complete the following diagram, which interprets all the ribbon numbers mentioned earlier.

Finally we consider a family of quotient rings $R_\mu = \mathbb{F}[X]/I_\mu$ indexed by partitions μ of n , which contains the coinvariant algebra of $\mathfrak{S}_n$ as a special case ($\mu = 1^n$). If $\mathbb{F} = \mathbb{C}$ then R_μ is isomorphic to the cohomology rings of the Springer fiber $\mathcal{F}_\mu$, carries an $\mathfrak{S}_n$ -action, and has graded Frobenius characteristic equal to the modified Hall–Littlewood symmetric function $\tilde{H}_\mu(x; t)$. We prove that the $H_n(0)$ -action on $\mathbb{F}[X]$ preserves the ideal I_μ if and only if μ is a hook, and if so then R_μ becomes a projective $H_n(0)$ -module whose graded characteristic also equals $\tilde{H}_\mu(x; t)$, and whose graded non-commutative characteristic equals $\tilde{\mathbf{H}}_\mu(X; t)$, a non-commutative analogue of $\tilde{H}_\mu(x; t)$ introduced by Bergeron and Zabrocki [4].

This paper is structured as follows. First we review the definitions for the various ribbon numbers and their interpretation by representations of $\mathfrak{S}_n$ and $GL(n, \mathbb{F}_q)$ in Sect. 2. Then we recall the representation theory of the 0-Hecke algebra in Sect. 3. The result on the coinvariant algebra $\mathbb{F}[X]/(\mathbb{F}[X]_+^{\mathfrak{S}_n})$ is given in Sect. 4, and a generalization to the Laurent ring $\mathbb{F}[\Lambda]$ of the weight lattice Λ of a Weyl group is provided in Sect. 5. Next we investigate the 0-Hecke algebra actions on the flag variety 1_B^G and the coinvariant algebra of (G, B) in Sect. 6 and Sect. 7. The $H_n(0)$ -action on the cohomology rings of the Springer fibers is studied in Sect. 8. Lastly Sect. 9 is devoted to questions for future research.

2 Ribbon numbers

A *composition* is a sequence $\alpha = (\alpha_1, \dots, \alpha_\ell)$ of positive integers $\alpha_1, \dots, \alpha_\ell$. The *length* of α is $\ell(\alpha) := \ell$ and the *size* of α is $|\alpha| := \alpha_1 + \dots + \alpha_\ell$. If the size of α is n then say α is a composition of n and write $\alpha \models n$. Let $\sigma_j = \alpha_1 + \dots + \alpha_j$ for $j = 0, 1, \dots, \ell$; in particular, $\sigma_0 = 0$ and $\sigma_\ell = n$. The *descent set* of α is $D(\alpha) := \{\sigma_1, \dots, \sigma_{\ell-1}\}$. The map $\alpha \mapsto D(\alpha)$ is a bijection between compositions of n and subsets of $[n-1]$, where $[n] := \{1, \dots, n\}$. Write $\tilde{\alpha} := (\alpha_\ell, \dots, \alpha_1)$ and let α^c be the composition of n with $D(\alpha^c) = [n-1] \setminus D(\alpha)$. Write $\alpha \preceq \beta$ if α and β are two compositions of the same size with $D(\alpha) \subseteq D(\beta)$.

A *semistandard Young tableau* τ of an arbitrary skew shape λ/μ is a filling of the skew diagram of λ/μ by positive integers such that every row weakly increases from left to right and every column strictly increases from top to bottom. Reading these integers from the bottom row to the top row and proceeding from left to right within each row gives the *reading word* $w(\tau)$ of τ . Say τ is a *standard Young tableau* if the integers appearing in τ are precisely $1, \dots, n$ without repetition, i.e. $w(\tau) \in \mathfrak{S}_n$. The *descents* of a standard Young tableau τ are those numbers i appearing in a row strictly below $i+1$ in τ , or in other words, the descents of $w(\tau)^{-1}$. The *major index* $\text{maj}(\tau)$ of a standard Young tableau τ is the sum of all its descents. Denote by $\text{SSYT}(\lambda/\mu)$ [$\text{SYT}(\lambda/\mu)$ resp.] the set of all semistandard [standard resp.] Young tableaux of shape λ/μ .

A *ribbon* is a skew connected diagram without 2×2 boxes. A ribbon α whose rows have lengths $\alpha_1, \dots, \alpha_\ell$, ordered *from bottom to top*, can be identified with a composition $\alpha = (\alpha_1, \dots, \alpha_\ell)$. Denote by $\alpha^\sim$ the *transpose* of the ribbon α . One can

check that $\alpha^\sim = (\overleftarrow{\alpha})^c = \overleftarrow{\alpha^c}$. An example is given below.

A (standard) ribbon tableau is a standard Young tableau of ribbon shape α . Taking the reading word $\tau \mapsto w(\tau)$ gives a bijection between $\text{SYT}(\alpha)$ and the descent class of α , which consists of all permutations w in $\mathfrak{S}_n$ with $D(w) = D(\alpha)$; see e.g. Kroh and Thibon [21]. The descent class of α is an interval under the left weak order of $\mathfrak{S}_n$, denoted by $[w_0(\alpha), w_1(\alpha)]$. For instance, the descent class of $\alpha = (1, 2, 1)$ is given below.

In particular, the ribbon tableaux of $w_0(\alpha)$ and $w_1(\alpha)$ can be, respectively, obtained by

- filling with $1, 2, \dots, n$ the columns of the ribbon α from top to bottom, starting with the leftmost column and proceeding toward the rightmost column,
- filling with $1, 2, \dots, n$ the rows of the ribbon α from left to right, starting with the top row and proceeding toward the bottom row.

The ribbon number r_α is the cardinality of the descent class of α . The q -ribbon number is

$$r_\alpha(q) := \sum_{\substack{w \in \mathfrak{S}_n: \\ D(w) = D(\alpha)}} q^{\text{inv}(w)} = [n]!_q \det \left(\frac{1}{[\sigma_j - \sigma_{i-1}]!_q} \right)_{i,j=1}^\ell \quad (3)$$

where $[n]!_q = [n]_q [n-1]_q \cdots [1]_q$ and $[n]_q = 1 + q + \cdots + q^{n-1}$. By Foata and Schützenberger [11], inv and maj are equidistributed on every inverse descent class

$\{w \in \mathfrak{S}_n : D(w^{-1}) = D(\alpha)\}$. Thus

$$r_\alpha(t) = \sum_{\substack{w \in \mathfrak{S}_n : \\ D(w) = D(\alpha)}} t^{\text{maj}(w^{-1})} \xrightarrow{w(\tau) \leftrightarrow \tau} \sum_{\tau \in \text{SYT}(\alpha)} t^{\text{maj}(\tau)}.$$

A further generalization, introduced by Reiner and Stanton [27], is the (q, t) -ribbon number

$$r_\alpha(q, t) := \sum_{\substack{w \in \mathfrak{S}_n : \\ D(w) = D(\alpha)}} \text{wt}(w; q, t) = n!_{q,t} \det \left(\varphi^{\sigma_i - 1} \frac{1}{(\sigma_j - \sigma_{i-1})!_{q,t}} \right)_{i,j=1}^\ell.$$

Here $\text{wt}(w; q, t)$ is some weight defined by product expression, $n!_{q,t} = (1 - t^{q^m - 1})(1 - t^{q^m - q}) \cdots (1 - t^{q^m - q^{m-1}})$, and $\varphi : t \mapsto t^q$ is the Frobenius operator.

All these ribbon numbers can be interpreted by the homology representation χ^α [χ_q^α resp.] of $\mathfrak{S}_n$ [$G = GL(n, \mathbb{F}_q)$ resp.], defined as the top homology of the rank-selected Coxeter complex $\Delta(\mathfrak{S}_n)_\alpha$ [Tits building $\Delta(G)_\alpha$ resp.], and by the intertwiner $M^\alpha = \text{Hom}_{\mathbb{F}\mathfrak{S}_n}(\chi^\alpha, \mathbb{F}[X])$ [$M_q^\alpha = \text{Hom}_{\mathbb{F}G}(\chi_q^\alpha, \mathbb{F}[X])$ resp.] as a module over $\mathbb{F}[X]^{\mathfrak{S}_n}$ [$\mathbb{F}[X]^G$ resp.]. By work of Reiner and Stanton [27], one has the following picture.

The ribbon numbers are related to the multinomial coefficients by inclusion–exclusion. Let $\alpha = (\alpha_1, \dots, \alpha_\ell)$ be a composition of n . Then we have the *multinomial* and *q-multinomial coefficients*

$$\begin{aligned} \left[\begin{matrix} n \\ \alpha \end{matrix} \right] &:= \frac{n!}{\alpha_1! \cdots \alpha_\ell!} = \#\{w \in \mathfrak{S}_n : D(w) \subseteq D(\alpha)\}, \\ \left[\begin{matrix} n \\ \alpha \end{matrix} \right]_q &:= \frac{[n]!_q}{[\alpha_1]!_q \cdots [\alpha_\ell]!_q} = \sum_{\substack{w \in \mathfrak{S}_n : \\ D(w) \subseteq D(\alpha)}} q^{\text{inv}(w)}. \end{aligned}$$

Reiner and Stanton [27] introduced the (q, t) -multinomial coefficient

$$\left[\begin{matrix} n \\ \alpha \end{matrix} \right]_{q,t} := \frac{n!_{q,t}}{\alpha_1!_{q,t} \cdot \varphi^{\sigma_1}(\alpha_2!_{q,t}) \cdots \varphi^{\sigma_{\ell-1}}(\alpha_{\ell}!_{q,t})} = \sum_{w \in \mathfrak{S}_n : D(w) \subseteq D(\alpha)} \text{wt}(w; q, t).$$

Assume q is a prime power below. Let $G = GL(n, \mathbb{F}_q)$ be the finite general linear group over $\mathbb{F}_q$, and let P_{α} be the parabolic subgroup of all invertible block upper triangular matrices whose diagonal blocks have sizes given by the composition α . Then

$$\begin{aligned} \left[\begin{matrix} n \\ \alpha \end{matrix} \right]_q &= |G/P_{\alpha}|, \\ \left[\begin{matrix} n \\ \alpha \end{matrix} \right]_{q,t} &= \text{Hilb}(\mathbb{F}[X]^{P_{\alpha}} / (\mathbb{F}[X]_+^G), t). \end{aligned}$$

3 Representation theory of the 0-Hecke algebras

In this section we recall from Norton [26] and Krob and Thibon [21] the representation theory of the 0-Hecke algebras of finite Coxeter groups.

3.1 Simple modules and projective indecomposable modules

Let

$$W = \langle s_1, \dots, s_r : s_i^2 = 1, (s_i s_j s_i \cdots)_{m_{ij}} = (s_j s_i s_j \cdots)_{m_{ij}}, 1 \leq i \neq j \leq r \rangle$$

be a finite Coxeter group, where $(aba \cdots)_m$ denotes an alternating product of m terms. The 0-Hecke algebra $H_W(0)$ of W is an associative $\mathbb{F}$ -algebra generated by $T_1, \dots, T_r$ with relations

$$\begin{cases} T_i^2 = -T_i, & 1 \leq i \leq r, \\ (T_i T_j T_i \cdots)_{m_{ij}} = (T_j T_i T_j \cdots)_{m_{ij}}, & 1 \leq i \neq j \leq r. \end{cases} \quad (4)$$

Another set of generators $\{T'_i = T_i + 1 : i = 1, \dots, r\}$ for $H_W(0)$ satisfies the relations

$$\begin{cases} (T'_i)^2 = T'_i, & 1 \leq i \leq r, \\ (T'_i T'_j T'_i \cdots)_{m_{ij}} = (T'_j T'_i T'_j \cdots)_{m_{ij}}, & 1 \leq i \neq j \leq r. \end{cases} \quad (5)$$

Thus $T_w = T_{i_1} \cdots T_{i_k}$ and $T'_w = T'_{i_1} \cdots T'_{i_k}$ are both well-defined if $w = s_{i_1} \cdots s_{i_k}$ is a reduced expression.

Let α be a composition of $r + 1$. Similarly to the type A case (i.e. $W = \mathfrak{S}_n$) mentioned in the previous section, the *descent class* of α in W consists of all elements $w \in W$ with descent set

$$D(w) := \{s_i \in S : \ell(ws_i) < \ell(w)\} = \{s_i : i \in D(\alpha)\}.$$

It is an interval in the left weak order of W , denoted by $[w_0(\alpha), w_1(\alpha)]$, where $w_0(\alpha)$ is the longest element in the parabolic subgroup $W_{D(\alpha)}$; see Björner and Wachs [5, Theorem 6.2].

Norton [26] decomposed $H_W(0)$ into a direct sum of 2^r non-isomorphic indecomposable submodules

$$\mathbf{P}_\alpha := H_W(0) \cdot T_{w_0(\alpha)} T'_{w_0(\alpha^c)}$$

indexed by compositions α of $r + 1$; each $\mathbf{P}_\alpha$ has an $\mathbb{F}$ -basis

$$\{T_w T'_{w_0(\alpha^c)} : w \in [w_0(\alpha), w_1(\alpha)]\}.$$

The *top* of $\mathbf{P}_\alpha$, denoted by $\mathbf{C}_\alpha := \text{top}(\mathbf{P}_\alpha) = \mathbf{P}_\alpha / \text{rad } \mathbf{P}_\alpha$, is a (one-dimensional) simple $H_W(0)$ -module with the action of $H_W(0)$ given by

$$T_i = \begin{cases} -1, & \text{if } i \in D(\alpha), \\ 0, & \text{if } i \notin D(\alpha). \end{cases}$$

It follows from the general theory (see e.g. [2, Sect. I.5]) that $\{\mathbf{P}_\alpha : \alpha \models r + 1\}$ is a complete list of non-isomorphic projective indecomposable $H_W(0)$ -modules and $\{\mathbf{C}_\alpha : \alpha \models r + 1\}$ is a complete list of non-isomorphic simple $H_W(0)$ -modules.

3.2 0-Hecke algebras of the symmetric groups

Now assume we are in type A , i.e. $W = \mathfrak{S}_n$ and $H_W(0) = H_n(0)$. The projective indecomposable $H_n(0)$ -modules can be described in a combinatorial way [17, 21] using ribbon tableaux. We see in the previous section that the ribbon tableaux of shape α are in bijection with the descent class of α , hence in bijection with the basis of $\mathbf{P}_\alpha$ given above. The $H_n(0)$ -action on $\mathbf{P}_\alpha$ agrees with the following $H_n(0)$ -action on these ribbon tableaux:

$$T_i \tau = \begin{cases} -\tau, & \text{if } i \text{ is in a higher row of } \tau \text{ than } i + 1, \\ 0, & \text{if } i \text{ is in the same row of } \tau \text{ as } i + 1, \\ s_i \tau, & \text{if } i \text{ is in a lower row of } \tau \text{ than } i + 1, \end{cases} \quad (6)$$

where τ is a ribbon tableau of shape α and $s_i \tau$ is obtained from τ by swapping i and $i + 1$. This action gives rise to a directed version of the Hasse diagram of the interval $[w_0(\alpha), w_1(\alpha)]$ under the weak order. The top and bottom tableaux in this diagram correspond to $\mathbf{C}_\alpha = \text{top}(\mathbf{P}_\alpha)$ and $\mathbf{C}_{\bar{\alpha}} = \text{soc}(\mathbf{P}_\alpha)$. An example is given below for

$\alpha = (1, 2, 1)$:

3.3 Quasi-symmetric and non-commutative symmetric functions

Krob and Thibon [21] provided a correspondence between representations of $H_n(0)$ and the dual Hopf algebras **QSym** of quasi-symmetric functions and **NSym** of non-commutative symmetric functions.

The Hopf algebra **QSym** is a free $\mathbb{Z}$ -module on the *monomial quasi-symmetric functions*

$$M_\alpha := \sum_{1 \leq i_1 < \dots < i_\ell} x_{i_1}^{\alpha_1} \cdots x_{i_\ell}^{\alpha_\ell}$$

for all compositions $\alpha = (\alpha_1, \dots, \alpha_\ell)$. Another free $\mathbb{Z}$ -basis consists of the *fundamental quasi-symmetric functions*

$$F_\alpha := \sum_{\alpha \preceq \beta} M_\beta = \sum_{\substack{1 \leq i_1 \leq \dots \leq i_k \\ j \in D(\alpha) \Rightarrow i_j < i_{j+1}}} x_{i_1} \cdots x_{i_k}.$$

The Hopf algebra **NSym** is the free associative algebra $\mathbb{Z}\langle \mathbf{h}_1, \mathbf{h}_2, \dots \rangle$ where

$$\mathbf{h}_k := \sum_{1 \leq i_1 \leq \dots \leq i_k} X_{i_1} \cdots X_{i_k}, \quad \forall k \geq 1.$$

A free $\mathbb{Z}$ -basis for **NSym** consists of the *complete homogeneous non-commutative symmetric functions* $\mathbf{h}_\alpha := \mathbf{h}_{\alpha_1} \cdots \mathbf{h}_{\alpha_\ell}$ for all compositions $\alpha = (\alpha_1, \dots, \alpha_\ell)$. Another free $\mathbb{Z}$ -basis consists of the *non-commutative ribbon Schur functions*

$$\mathbf{s}_\alpha := \sum_{\beta \preceq \alpha} (-1)^{\ell(\alpha) - \ell(\beta)} \mathbf{h}_\beta$$

for all compositions α . The duality between $\mathbf{QSym}$ and $\mathbf{NSym}$ is given by $\langle M_\alpha, \mathbf{h}_\beta \rangle = \langle F_\alpha, \mathbf{s}_\beta \rangle := \delta_{\alpha\beta}$.

The dual Hopf algebras $\mathbf{QSym}$ and $\mathbf{NSym}$ are also related to the self-dual Hopf algebra $\mathbf{Sym}$, the *ring of symmetric functions*. A *positive self-dual basis* for $\mathbf{Sym}$ consists of the *Schur functions* s_λ for all partitions λ . The definition of s_λ is a special case of the *skew Schur function*

$$s_{\lambda/\mu} := \sum_{\tau \in \text{SSYT}(\lambda/\mu)} x^\tau$$

of a skew shape λ/μ , where $x^\tau := x_1^{d_1} x_2^{d_2} \cdots$ if τ is of type $(d_1, d_2, \dots)$, i.e. $d_1, d_2, \dots$ are the multiplicities of $1, 2, \dots$ in τ . The commutative image of a non-commutative ribbon Schur function $\mathbf{s}_\alpha$ is nothing but the ribbon Schur function s_α . This gives a surjection $\mathbf{NSym} \twoheadrightarrow \mathbf{Sym}$.

There is also a free $\mathbb{Z}$ -basis for $\mathbf{Sym}$ consisting of the *monomial symmetric functions*

$$m_\lambda := \sum_{\lambda(\alpha)=\lambda} M_\alpha$$

for all partitions λ . Here $\lambda(\alpha)$ is the unique partition obtained from the composition α by rearranging its parts. This gives an injection (actually an inclusion) $\mathbf{Sym} \hookrightarrow \mathbf{QSym}$.

We will use the following expansion of a Schur function indexed by a partition $\lambda \vdash n$:

$$s_\lambda = \sum_{\mu \vdash n} K_{\lambda\mu} m_\mu. \quad (7)$$

Here $K_{\lambda\mu}$ is the Kostka number which counts all semistandard Young tableaux of shape λ and type μ .

3.4 Characteristic maps

Let A be an $\mathbb{F}$ -algebra and let $\mathcal{C}$ be a category of some finitely generated A -modules. The *Grothendieck group* of $\mathcal{C}$ is defined as the abelian group F/R , where F is the free abelian group on the isomorphism classes $[M]$ of the $H_n(0)$ -modules M in $\mathcal{C}$, and R is the subgroup of F generated by the elements $[M] - [L] - [N]$ corresponding to all exact sequences $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ of A -modules in $\mathcal{C}$. Note that every exact sequence of projective modules splits.

Denote by $G_0(\mathfrak{S}_n)$ the Grothendieck group of category of all finitely generated $\mathbb{C}\mathfrak{S}_n$ -modules. Then $G_0(\mathfrak{S}_n)$ is a free abelian group on the isomorphism classes of simple $\mathbb{C}\mathfrak{S}_n$ -modules $[S_\lambda]$ for all $\lambda \vdash n$. The tower of groups $\mathfrak{S}_\bullet : \mathfrak{S}_0 \hookrightarrow \mathfrak{S}_1 \hookrightarrow \mathfrak{S}_2 \hookrightarrow \cdots$ has a Grothendieck group

$$G_0(\mathfrak{S}_\bullet) := \bigoplus_{n \geq 0} G_0(\mathfrak{S}_n).$$

This is a self-dual Hopf algebra with product and coproduct given by induction and restriction of representations along the natural embeddings $\mathfrak{S}_m \times \mathfrak{S}_n \hookrightarrow \mathfrak{S}_{m+n}$. The

Frobenius characteristic map ch sends a simple S_λ to the Schur function s_λ , giving a Hopf algebra isomorphism $G_0(\mathfrak{S}_\bullet) \cong \text{Sym}$.

There is an analogous result for $H_n(0)$ -modules (over an arbitrary field $\mathbb{F}$). The Grothendieck group of the category of all finitely generated $H_n(0)$ -modules is denoted by $G_0(H_n(0))$, and the Grothendieck group of the category of finitely generated projective $H_n(0)$ -modules is denoted by $K_0(H_n(0))$. We have

$$G_0(H_n(0)) = \bigoplus_{\alpha \models n} \mathbb{Z} \cdot [\mathbf{C}_\alpha], \quad K_0(H_n(0)) = \bigoplus_{\alpha \models n} \mathbb{Z} \cdot [\mathbf{P}_\alpha].$$

The two Grothendieck groups of the tower of algebras $H_\bullet(0) : H_0(0) \hookrightarrow H_1(0) \hookrightarrow H_2(0) \hookrightarrow \dots$ are

$$G_0(H_\bullet(0)) := \bigoplus_{n \geq 0} G_0(H_n(0)), \quad K_0(H_\bullet(0)) := \bigoplus_{n \geq 0} K_0(H_n(0)).$$

These two Grothendieck groups are dual Hopf algebras with product and coproduct again given by induction and restriction of representations along the embeddings $H_m(0) \otimes H_n(0) \hookrightarrow H_{m+n}(0)$. Krob and Thibon [21] defined two Hopf algebra isomorphisms:

$$\text{Ch} : G_0(H_\bullet(0)) \cong \text{QSym}, \quad \mathbf{ch} : K_0(H_\bullet(0)) \cong \text{NSym}.$$

The *quasi-symmetric characteristic map* Ch sends a finitely generated $H_n(0)$ -module M with simple composition factors $\mathbf{C}_{\alpha(1)}, \dots, \mathbf{C}_{\alpha(k)}$ to

$$\text{Ch}(M) := F_{\alpha(1)} + \dots + F_{\alpha(k)}.$$

Similarly, the *non-commutative characteristic map* $\mathbf{ch}$ sends a finitely generated projective $H_n(0)$ -module $M = \mathbf{P}_{\alpha(1)} \oplus \dots \oplus \mathbf{P}_{\alpha(k)}$ to

$$\mathbf{ch}(M) := \mathbf{s}_{\alpha(1)} + \dots + \mathbf{s}_{\alpha(k)}.$$

Krob and Thibon [21] also showed that $\text{Ch}(\mathbf{P}_\alpha) = s_\alpha$, which is the commutative image of $\mathbf{ch}(\mathbf{P}_\alpha) = \mathbf{s}_\alpha$. Thus $\text{Ch}(M)$ is symmetric whenever M is a finitely generated projective $H_n(0)$ -module, but not vice versa (e.g. $\mathbf{C}_{12} \oplus \mathbf{C}_{21}$ is nonprojective and has quasi-symmetric characteristic equal to s_{21}).

If $M = H_n(0)v$ is cyclic then the *length filtration*

$$H_n(0)^{(\ell)} = \bigoplus_{\ell(w) \geq \ell} \mathbb{F} T_w$$

induces a filtration of $H_n(0)$ -modules $M^{(\ell)} = H_n(0)^{(\ell)}v$ for all $\ell \geq 0$. This refines $\text{Ch}(M)$ to a *graded characteristic*

$$\text{Ch}_q(M) = \sum_{\ell \geq 0} q^\ell \text{Ch}(M^{(\ell)} / M^{(\ell+1)}).$$

Taking M to be the regular representation of $H_n(0)$ we have

$$\mathrm{Ch}_q(H_n(0)) = \sum_{w \in \mathfrak{S}_n} q^{\mathrm{inv}(w)} F_{D(w^{-1})} = \sum_{\alpha} r_{\alpha}(q) F_{\alpha}$$

and taking a limit as $q \rightarrow 1$ gives

$$\mathrm{Ch}(H_n(0)) = \sum_{\alpha} r_{\alpha} F_{\alpha}.$$

If M has another filtration by $H_n(0)$ -modules M_d for $d \geq 0$, then one can look at the bifiltration of $H_n(0)$ -modules $M^{(\ell, d)} = M^{(\ell)} \cap M_d$ for $\ell, d \geq 0$, and define the *bigraded characteristic* to be

$$\mathrm{Ch}_{q,t}(M) = \sum_{\ell, d \geq 0} q^{\ell} t^d \mathrm{Ch}(M^{(\ell, d)} / (M^{(\ell+1, d)} + M^{(\ell, d+1)})).$$

If a projective $H_n(0)$ -module $M = \mathbf{P}_{\alpha^{(1)}} \oplus \cdots \oplus \mathbf{P}_{\alpha^{(k)}}$ has a grading $\deg \mathbf{P}_{\alpha^{(i)}} = d_i$ then we can define the graded non-commutative characteristic of M to be

$$\mathbf{ch}_t(M) := t^{d_1} \mathbf{s}_{\alpha^{(1)}} + \cdots + t^{d_k} \mathbf{s}_{\alpha^{(k)}}.$$

4 Coinvariant algebra of $H_n(0)$

In this section we give interpretations of the ribbon and q -ribbon numbers by studying the $H_n(0)$ -action on its coinvariant algebra.

The symmetric group $\mathfrak{S}_n$ acts on the polynomial ring $\mathbb{F}[X] := \mathbb{F}[x_1, \dots, x_n]$ over an arbitrary field $\mathbb{F}$ by permuting the variables $x_1, \dots, x_n$, and hence acts on the coinvariant algebra $\mathbb{F}[X]/(\mathbb{F}[X]_{+}^{\mathfrak{S}_n})$ of $\mathfrak{S}_n$, where $(\mathbb{F}[X]_{+}^{\mathfrak{S}_n})$ is the ideal generated by symmetric polynomials of positive degree. We often identify the polynomials in $\mathbb{F}[X]$ with their images in the coinvariant algebra of $\mathfrak{S}_n$ in this section.

For $i = 1, \dots, n-1$, let s_i is the adjacent transposition $(i, i+1)$, and define the *Demazure operators*

$$\pi_i f := \frac{x_i f - x_{i+1} s_i(f)}{x_i - x_{i+1}}, \quad \forall f \in \mathbb{F}[X]. \quad (8)$$

It follows from this definition that

- π_i satisfy the same relations as T'_i , i.e. the relations in (5),
- $\deg(\pi_i f) = \deg(f)$ for all homogeneous polynomials $f \in \mathbb{F}[X]$,
- $\pi_i f = f$ if and only if $s_i f = f$ for all $f \in \mathbb{F}[X]$,
- $\pi_i(fg) = f\pi_i(g)$ for all $f, g \in \mathbb{F}[X]$ satisfying $\pi_i f = f$.

Hence we have an $H_n(0)$ -action on $\mathbb{F}[X]$ by $T'_i \mapsto \pi_i$, or equivalently $T_i \mapsto \bar{\pi}_i := \pi_i - 1$, for $i = 1, \dots, n-1$. The operators π_w and $\bar{\pi}_w$ are defined similarly as T'_w

and T_w for all $w \in \mathfrak{S}_n$. This $H_n(0)$ -action preserves the degrees, has the same invariants as $\mathfrak{S}_n$, and is $\mathbb{F}[X]^{\mathfrak{S}_n}$ -linear. Thus we have an $H_n(0)$ -action on the *coinvariant algebra of $H_n(0)$* , which is defined as

$$\mathbb{F}[X]/(f \in \mathbb{F}[X] : \deg f > 0, \pi_i f = f, 1 \leq i \leq n-1)$$

and coincides with the coinvariant algebra of $\mathfrak{S}_n$.

To study the $H_n(0)$ -action on its coinvariant algebra, we consider certain *Demazure atoms* which behave nicely under the $H_n(0)$ -action, i.e. the polynomials $\pi_w x_{D(w)}$ for all $w \in \mathfrak{S}_n$. Here

$$x_I := \prod_{i \in I} x_1 \cdots x_i.$$

for any $I \subseteq [n-1]$. See Mason [25] for more information on the Demazure atoms.

We will see in Lemma 4.1 that the Demazure atoms mentioned above are closely related to the *descent monomials*

$$w x_{D(w)} = \prod_{i \in D(w)} x_{w(1)} \cdots x_{w(i)}, \quad \forall w \in \mathfrak{S}_n.$$

It is well known that the descent monomials form a basis for the coinvariant algebra of $\mathfrak{S}_n$; see e.g. Garsia [12] and Steinberg [31]. Allen [1] provided an elementary proof for this result, which we will adapt to the Demazure atoms $\pi_w x_{D(w)}$. Thus we first recall Allen's proof below.

A *weak composition* is a finite sequence of nonnegative integers. A *partition* is a finite decreasing sequence of nonnegative integers, with zeros ignored sometimes. Every monomial in $\mathbb{F}[X]$ can be written as $x^d = x_1^{d_1} \cdots x_n^{d_n}$ where $d = (d_1, \dots, d_n)$ is a weak composition. Denote by $\lambda(d)$ the unique partition obtained from rearranging the weak composition d . Given two monomials x^d and x^e , write $x^d < x^e$ or $d < e$ if $\lambda(d) <_L \lambda(e)$, and write $x^d <_{\text{ts}} x^e$ if (i) $\lambda(d) <_L \lambda(e)$ or (ii) $\lambda(d) = \lambda(e)$ and $d <_L e$, where “ $<_L$ ” is the lexicographic order.

Given a weak composition $d = (d_1, \dots, d_n)$, we have a permutation $\sigma(d) \in \mathfrak{S}_n$ obtained by labeling $d_1, \dots, d_n$ from the largest to the smallest, breaking ties from left to right. Construct a weak composition $\gamma(d)$ from this labeling as follows. First replace the largest label with 0, and recursively, if the label t has been replaced with s , then replace $t-1$ with s if it is to the left of t , or with $s+1$ otherwise. Let $\mu(d) = d - \gamma(d)$ be the component-wise difference. For example,

$$\begin{aligned} d &= (3, 1, 3, 0, 2, 0), & \sigma(d) &= (1, 4, 2, 5, 3, 6), \\ \gamma(d) &= (1, 0, 1, 0, 1, 0), & \mu(d) &= (2, 1, 2, 0, 1, 0). \end{aligned}$$

The decomposition $d = \gamma(d) + \mu(d)$ is the usual *P-partition encoding* of d (see e.g. Stanley [30]), and $x^{\gamma(d)}$ is the descent monomial of $\sigma(d)^{-1}$. E.E. Allen [1, Proposition 2.1] showed that $w\mu(d) + \gamma(d) <_{\text{ts}} d$ for all $w \in \mathfrak{S}_n$ unless $w = 1$, and thus

$$m_{\mu(d)} \cdot x^{\gamma(d)} = x^d + \sum_{x^e <_{\text{ts}} x^d} c_e x^e, \quad c_e \in \mathbb{Z}, \quad (9)$$

where $m_{\mu(d)}$ is the monomial symmetric function corresponding to $\mu(d)$, i.e. the sum of the monomials in the $\mathfrak{S}_n$ -orbit of $x^{\mu(d)}$. It follows that

$$\{m_{\mu} \cdot wx_{D(w)} : \mu = (\mu_1 \geq \cdots \geq \mu_n \geq 0), w \in \mathfrak{S}_n\}$$

is triangularly related to the set of all monomials x^d , and thus an $\mathbb{F}$ -basis for $\mathbb{F}[X]$. Therefore the descent monomials form an $\mathbb{F}[X]^{\mathfrak{S}_n}$ -basis for $\mathbb{F}[X]$ and give an $\mathbb{F}$ -basis for $\mathbb{F}[X]/(\mathbb{F}[X]_+^{\mathfrak{S}_n})$.

Now we investigate the relation between our Demazure atoms and the descent monomials. First observe that if m is a monomial not containing x_i and x_{i+1} , then

$$\bar{\pi}_i(mx_i^a x_{i+1}^b) = \begin{cases} m(x_i^{a-1} x_{i+1}^{b+1} + x_i^{a-2} x_{i+1}^{b+2} + \cdots + x_i^b x_{i+1}^a), & \text{if } a > b, \\ 0, & \text{if } a = b, \\ -m(x_i^a x_{i+1}^b - x_i^{a+1} x_{i+1}^{b-1} - \cdots - x_i^{b-1} x_{i+1}^{a+1}), & \text{if } a < b. \end{cases} \quad (10)$$

Lemma 4.1 *Suppose that α is a composition of n and w is a permutation in $\mathfrak{S}_n$ with $D(w) \subseteq D(\alpha)$. Then*

$$\bar{\pi}_{wx_{D(\alpha)}} = wx_{D(\alpha)} + \sum_{x^d \prec x_{D(\alpha)}} c_d x^d, \quad c_d \in \mathbb{Z}.$$

Moreover, $wx_{D(\alpha)}$ is a descent monomial if and only if $D(w) = D(\alpha)$.

Proof We prove the first assertion by induction on $\ell(w)$. If $\ell(w) = 0$ then we are done; otherwise we assume $w = s_j u$ for some $j \in [n-1]$ and some $u \in \mathfrak{S}_n$ with $\ell(u) < \ell(w)$. Since $D(u) \subseteq D(w) \subseteq D(\alpha)$, one has

$$\bar{\pi}_{ux_{D(\alpha)}} = ux_{D(\alpha)} + \sum_{x^d \prec x_{D(\alpha)}} c_d x^d, \quad c_d \in \mathbb{Z}. \quad (11)$$

It follows from (10) that

$$\bar{\pi}_j(x^d) = \sum_{x^e \preceq x^d} a_e x^e, \quad a_e \in \mathbb{Z}. \quad (12)$$

Observe that the degree of x_k in $ux_{D(\alpha)}$ is

$$r_k = \#\{i \in D(\alpha) : u^{-1}(k) \leq i\}.$$

Since $\ell(s_j u) > \ell(u)$, we have $u^{-1}(j) < u^{-1}(j+1)$ and thus $r_j \geq r_{j+1}$. Since $(s_j u)^{-1}(j+1) < (s_j u)^{-1}(j)$, there exists an $i \in D(s_j u) \subseteq D(\alpha)$ such that

$$u^{-1}(j) = (s_j u)^{-1}(j+1) \leq i < (s_j u)^{-1}(j) = u^{-1}(j+1).$$

Thus $r_j > r_{j+1}$. It follows from (10) that

$$\bar{\pi}_j(ux_{D(\alpha)}) = s_j ux_{D(\alpha)} + \sum_{x^e \prec x_{D(\alpha)}} b_e x^e, \quad b_e \in \mathbb{Z}. \quad (13)$$

Combining (11), (12), and (13) proves the first assertion.

If $D(w) = D(\alpha)$ then $w x_{D(\alpha)}$ is the descent monomial of w . Conversely, assume $w x_{D(\alpha)}$ equals the descent monomial of some $u \in W$, i.e.

$$\prod_{i \in D(\alpha)} x_{w(1)} \cdots x_{w(i)} = \prod_{j \in D(u)} x_{u(1)} \cdots x_{u(j)}.$$

Let $D(\alpha) = \{i_1, \dots, i_k\}$ and $D(u) = \{j_1, \dots, j_t\}$. Comparing the variables absent on both sides of the above equality, one sees that $i_k = j_t$ and $w(i) = u(i)$ for $i = i_k + 1, \dots, n$. Repeat this argument for the variables appearing exactly m times, $m = 1, 2, \dots$, one sees that $D(\alpha) = D(u)$ and $w = u$. $\square$

Remark 4.2 Using the combinatorial formula by Mason [25] for the Demazure atoms, one can check that $\overline{\pi}_{w_0(\alpha) x_{D(\alpha)}}$ and $\overline{\pi}_{w_1(\alpha) x_{D(\alpha)}}$ are precisely the descent monomials of $w_0(\alpha)$ and $w_1(\alpha)$.

Lemma 4.3 For any weak composition $d = (d_1, \dots, d_n)$, let $\sigma = \sigma(d)$, $\gamma = \gamma(d)$, $\mu = \mu(d)$. If $c_\beta \in \mathbb{Z}$ for all $\beta \prec \gamma$ then

$$m_\mu \cdot \left(x^\gamma + \sum_{\beta \prec \gamma} c_\beta x^\beta \right) = x^d + \sum_{x^e \prec_{\text{ls}} x^d} b_e x^e, \quad b_e \in \mathbb{Z}.$$

Proof Since we already have (9), it suffices to show that $w\mu + \beta \prec d$ for all permutations w in $\mathfrak{S}_n$ and all $\beta \prec \gamma$. Given a weak composition α , let α_i be its i th part. Since $\sigma\mu$ and $\sigma\gamma$ are both weakly decreasing, one has $\lambda(\mu)_i + \lambda(\gamma)_i = \lambda(d)_i$ for all $i = 1, \dots, n$. Since $\beta \prec \gamma$, there exists a unique integer k such that $\lambda(\beta)_i = \lambda(\gamma)_i$ for $i = 1, \dots, k-1$, and $\lambda(\beta)_k < \lambda(\gamma)_k$. Then for all $i \in [k-1]$,

$$\begin{aligned} \lambda(w\mu + \beta)_i &\leq \lambda(\mu)_i + \lambda(\beta)_i \\ &= \lambda(\mu)_i + \lambda(\gamma)_i \\ &= \lambda(d)_i \end{aligned}$$

and

$$\begin{aligned} \lambda(w\mu + \beta)_k &\leq \lambda(\mu)_k + \lambda(\beta)_k \\ &< \lambda(\mu)_k + \lambda(\gamma)_k \\ &= \lambda(d)_k. \end{aligned}$$

Therefore $w\mu + \beta \prec d$ and we are done. $\square$

Lemma 4.4 The coinvariant algebra $\mathbb{F}[X]/(\mathbb{F}[X]_+^{\mathfrak{S}_n})$ has a basis given by $\{f_w : w \in \mathfrak{S}_n\}$, if

$$f_w = w x_{D(w)} + \sum_{x^d \prec_{X_{D(\alpha)}} c_d x^d, \quad c_d \in \mathbb{F}, \quad \forall w \in \mathfrak{S}_n.$$

Proof Given a weak composition $d = (d_1, \dots, d_n)$, let $\gamma = \gamma(d)$, $\mu = \mu(d)$, and $\sigma = \sigma(d)$. Then x^γ is the descent monomial of σ^{-1} . By Lemma 4.3,

$$m_\mu f_{\sigma^{-1}} = x^d + \sum_{x^e <_{\text{is}} x^d} b_e x^e.$$

Hence $\{m_\mu f_w : \mu = (\mu_1 \geq \dots \geq \mu_n \geq 0), w \in \mathfrak{S}_n\}$ is triangularly related to the set of all monomials x^d , and thus a basis for $\mathbb{F}[X]$. It follows that $\{f_w : w \in \mathfrak{S}_n\}$ is an $\mathbb{F}[X]^{\mathfrak{S}_n}$ -basis for $\mathbb{F}[X]$ and gives an $\mathbb{F}$ -basis for $\mathbb{F}[X]/(\mathbb{F}[X]_+^{\mathfrak{S}_n})$. $\square$

Theorem 4.5 *The coinvariant algebra $\mathbb{F}[X]/(\mathbb{F}[X]_+^{\mathfrak{S}_n})$ has an $\mathbb{F}$ -basis $\{\bar{\pi}_w x_{D(w)} : w \in \mathfrak{S}_n\}$ and decomposes as*

$$\mathbb{F}[X]/(\mathbb{F}[X]_+^{\mathfrak{S}_n}) = \bigoplus_{\alpha \models n} H_n(0) \cdot \bar{\pi}_{w_0(\alpha)} x_{D(\alpha)}$$

where $H_n(0) \cdot \bar{\pi}_{w_0(\alpha)} x_{D(\alpha)}$ has a basis $\{\bar{\pi}_w x_{D(\alpha)} : w \in [w_0(\alpha), w_1(\alpha)]\}$, and is isomorphic to the projective indecomposable $H_n(0)$ -module $\mathbf{P}_\alpha$, for all $\alpha \models n$. Consequently, $\mathbb{F}[X]/(\mathbb{F}[X]_+^{\mathfrak{S}_n})$ is isomorphic to the regular representation of $H_n(0)$.

Proof By Lemma 4.1 and Lemma 4.4, $\{\bar{\pi}_w x_{D(w)} : w \in \mathfrak{S}_n\}$ is a basis for $\mathbb{F}[X]/(\mathbb{F}[X]_+^{\mathfrak{S}_n})$. For any permutation u in $\mathfrak{S}_n$, one sees from the relations of $\bar{\pi}_i$ that $\bar{\pi}_u \bar{\pi}_{w_0(\alpha)} = \pm \bar{\pi}_w$ for some $w \geq w_0(\alpha)$ in the left weak order, which implies $D(w) \supseteq D(\alpha)$. If there exists $j \in D(w) \setminus D(\alpha)$, then

$$\bar{\pi}_w x_{D(\alpha)} = \bar{\pi}_{ws_j} \bar{\pi}_j x_{D(\alpha)} = 0.$$

Hence $H_n(0) \cdot \bar{\pi}_{w_0(\alpha)} x_{D(\alpha)}$ is spanned by $\{\bar{\pi}_w x_{D(\alpha)} : w \in [w_0(\alpha), w_1(\alpha)]\}$, which must be an $\mathbb{F}$ -basis since it is a subset of a linearly independent set. Sending $\bar{\pi}_w x_{D(\alpha)}$ to $T_w T'_{w_0(\alpha^c)}$ for all $w \in [w_0(\alpha), w_1(\alpha)]$ gives an isomorphism between $H_n(0) \cdot \bar{\pi}_{w_0(\alpha)} x_{D(\alpha)}$ and $\mathbf{P}_\alpha$. $\square$

Remark 4.6

- (i) This theorem and its proof are valid when $\mathbb{F}$ is replaced with $\mathbb{Z}$.
- (ii) By Remark 4.2, the cyclic generators $\bar{\pi}_{w_0(\alpha)} x_{D(\alpha)}$ for the indecomposable summands of the coinvariant algebra are precisely the descent monomials $w_0(\alpha) x_{D(\alpha)}$.

Theorem 4.7 (Lusztig–Stanley) *The graded Frobenius characteristic of the coinvariant algebra is*

$$\text{ch}_t(\mathbb{C}[X]/(\mathbb{C}[X]_+^{\mathfrak{S}_n})) = \sum_{\lambda \vdash n} \sum_{\tau \in \text{SYT}(\lambda)} t^{\text{maj}(\tau)} s_\lambda \stackrel{(*)}{=} \tilde{H}_{1^n}(x; t)$$

where $\tilde{H}_{1^n}(x; t)$ is the modified Hall–Littlewood symmetric function of the partition 1^n .

Remark 4.8 The equality (*) is a special case of Theorem 8.1. One can also see it by using the charge formula of Lascoux and Schützenberger [24].

We have an analogues result for the $H_n(0)$ -action on the coinvariant algebra.

Corollary 4.9

(i) *The bigraded characteristic of the coinvariant algebra is*

$$\mathrm{Ch}_{q,t}(\mathbb{F}[X]/(\mathbb{F}[X]_+^{\mathfrak{S}_n})) = \sum_{w \in \mathfrak{S}_n} t^{\mathrm{maj}(w)} q^{\mathrm{inv}(w)} F_{D(w^{-1})}$$

(ii) *The (degree) graded quasi-symmetric characteristic of the coinvariant algebra is*

$$\mathrm{Ch}_t(\mathbb{F}[X]/(\mathbb{F}[X]_+^{\mathfrak{S}_n})) = \sum_{\alpha \models n} r_\alpha(t) F_\alpha = \sum_{\alpha \models n} \sum_{\tau \in \mathrm{SYT}(\alpha)} t^{\mathrm{maj}(\tau)} F_\alpha.$$

(iii) *The quasi-symmetric function in (ii) is actually symmetric and equals*

$$\sum_{\lambda \vdash n} \left[\begin{matrix} n \\ \lambda \end{matrix} \right]_t m_\lambda = \sum_{\lambda \vdash n} \sum_{\tau \in \mathrm{SYT}(\lambda)} t^{\mathrm{maj}(\tau)} s_\lambda = \sum_{\lambda \vdash n} t^{n(\lambda)} \frac{[n]!_t}{\prod_{u \in \lambda} [h_u]_t} s_\lambda = \tilde{H}_{1^n}(x; t)$$

where h_u is the hook length of the box u in the Ferrers diagram of λ and $n(\lambda) := \lambda_2 + 2\lambda_3 + 3\lambda_4 + \dots$.

Proof Given a composition α of n , the $H_n(0)$ -module $H_n(0) \cdot \overline{\pi}_{w_0(\alpha)} x_{D(\alpha)}$ consists of homogeneous elements of degree $\mathrm{maj}(w)$ for any $w \in [w_0(\alpha), w_1(\alpha)]$. Hence Theorem 4.5 implies (i). It follows from (i) that the degree graded multiplicity of a simple $H_n(0)$ -module C_α in the coinvariant algebra is

$$r_\alpha(t) = \sum_{\substack{w \in \mathfrak{S}_n: \\ D(w^{-1}) = D(\alpha)}} t^{\mathrm{maj}(w)} = \sum_{\substack{w \in \mathfrak{S}_n: \\ D(w) = D(\alpha)}} t^{\mathrm{maj}(w^{-1})} = \sum_{\tau \in \mathrm{SYT}(\alpha)} t^{\mathrm{maj}(\tau)}.$$

Then

$$\mathrm{Ch}_t(\mathbb{F}[X]/(\mathbb{F}[X]_+^{\mathfrak{S}_n})) = \sum_{\alpha \models n} r_\alpha(t) F_\alpha = \sum_{\alpha \models n} \left[\begin{matrix} n \\ \alpha \end{matrix} \right]_t M_\alpha = \sum_{\lambda \vdash n} \left[\begin{matrix} n \\ \lambda \end{matrix} \right]_t m_\lambda \in \mathrm{Sym}[t].$$

Given a partition μ of n , we have

$$\left[\begin{matrix} n \\ \mu \end{matrix} \right]_t = \sum_{\substack{w \in \mathfrak{S}_n: \\ D(w) \subseteq D(\mu)}} t^{\mathrm{maj}(w^{-1})} = \sum_{w \in \mathfrak{S}(\mu)} t^{\mathrm{maj}(w^{-1})}$$

where $\mathfrak{S}(\mu)$ is the set of all permutations of the multiset of type μ . For example, $w = 3561247$ corresponds to

$$\begin{pmatrix} 1 & 1 & 1 & 2 & 2 & 3 & 3 \\ 3 & 5 & 6 & 1 & 2 & 4 & 7 \end{pmatrix} \in \mathfrak{S}(332).$$

Applying RSK to $w \in \mathfrak{S}(\mu)$ gives a pair (P, Q) of Young tableaux P and Q of the same shape (say λ), where P is standard, and Q is semistandard of type μ . It is well known that the descents of w^{-1} are precisely the descents of P ; see Schützenberger [28]. Hence

$$\begin{aligned} \text{Ch}_t(\mathbb{F}[X]/(\mathbb{F}[X]_+^{\mathfrak{S}_n})) &= \sum_{\lambda \vdash n} \sum_{P \in \text{SYT}(\lambda)} t^{\text{maj}(P)} \sum_{\mu \vdash n} K_{\lambda\mu} m_\mu \\ &= \sum_{\lambda \vdash n} t^{n(\lambda)} \frac{[n]_t!}{\prod_{u \in \lambda} [h_u]_t} s_\lambda. \end{aligned}$$

Here the last equality follows from the q -hook length formula and (7). $\square$

5 Coinvariant algebra of Weyl groups

The results in the previous section can be generalized to the action of the 0-Hecke algebra of a Weyl group W on the Laurent ring $\mathbb{F}[\Lambda]$ of the weight lattice Λ of W . The readers are referred to Humphreys [19] for details on the Weyl groups and weight theory.

Demazure's character formula [6] expresses the character of the highest weight modules over a semisimple Lie algebra using the *Demazure operators* π_i on the group ring $\mathbb{F}[\Lambda]$ of the weight lattice Λ . Write e^λ for the element in $\mathbb{F}[\Lambda]$ corresponding to the weight $\lambda \in \Lambda$. Suppose that $\gamma_1, \dots, \gamma_r$ are the simple roots,¹ $s_1, \dots, s_r$ are the simple reflections, and $\lambda_1, \dots, \lambda_r$ are the fundamental weights. Then

$$\mathbb{F}[\Lambda] = \mathbb{F}[z_1, \dots, z_r, z_1^{-1}, \dots, z_r^{-1}]$$

where $z_i = e^{\lambda_i}$. The *Demazure operators* are defined by

$$\pi_i = \frac{f - e^{-\gamma_i} s_i(f)}{1 - e^{-\gamma_i}}, \quad \forall f \in \mathbb{F}[\Lambda].$$

It follows that

$$\pi_i(e^\lambda) = \begin{cases} e^\lambda + e^{\lambda - \gamma_i} + \dots + e^{s_i \lambda}, & \text{if } \langle \lambda, \gamma_i \rangle \geq 0, \\ 0, & \text{if } \langle \lambda, \gamma_i \rangle = -1, \\ -e^{\lambda + \gamma_i} - \dots - e^{s_i \lambda - \gamma_i}, & \text{if } \langle \lambda, \gamma_i \rangle < -1. \end{cases} \quad (14)$$

Here $\langle \lambda, \gamma_i \rangle = 2(\lambda, \gamma_i)/(\gamma_i, \gamma_i)$ with $(-, -)$ being the standard inner product. See, for example, Kumar [23]. The Demazure operators satisfy $s_i \pi_i = \pi_i$, $\pi_i^2 = \pi_i$, and the braid relations [6, Sect. 5.5]. Hence the 0-Hecke algebra $H_W(0)$ of the Weyl group W acts on $\mathbb{F}[\Lambda]$ by sending T_i to $\pi_i = \pi_i - 1$. If $w = s_{i_1} \cdots s_{i_k}$ is a reduced expression then T_w acts by $\pi_w = \pi_{i_1} \cdots \pi_{i_k}$. It is clear that $\pi_i f = f$ if and only if $s_i f = s_i f$ for all $f \in \mathbb{F}[\Lambda]$.

¹We use γ to denote roots because α is used for compositions throughout this paper.

Using the Stanley–Reisner ring of the Coxeter complex of W , Garsia and Stanton [14] showed that

$$\mathbb{F}[\Lambda]^W = \mathbb{F}[a_1, \dots, a_r]$$

where

$$a_i = \sum_{w \in W/W_{i^c}} e^{w\lambda_i} \quad (i^c = [r] \setminus \{i\})$$

and $\mathbb{F}[\Lambda]$ has a free basis over $\mathbb{F}[\Lambda]^W$, which consists of the *descent monomials*

$$z_w := \prod_{i \in D(w)} e^{w\lambda_i}, \quad \forall w \in W.$$

See also Steinberg [31]. If we write $\lambda_I = \sum_{i \in I} \lambda_i$ for all subsets $I \subseteq [r]$, then $z_w = w e^{\lambda_{D(w)}}$. The basis $\{z_w : w \in W\}$ induces an $\mathbb{F}$ -basis for $\mathbb{F}[\Lambda]/(a_1, \dots, a_r)$. The $H_W(0)$ -action on $\mathbb{F}[\Lambda]$ is $\mathbb{F}[\Lambda]^W$ -linear, hence inducing an $H_W(0)$ -action on $\mathbb{F}[\Lambda]/(a_1, \dots, a_r)$.

We order the weights by $\lambda \leq \mu$ if $\mu - \lambda$ is a nonnegative linear combination of simple roots. Every monomial in $\mathbb{F}[\Lambda]$ is of the form $m = e^\lambda$ for some weight λ . By Humphreys [19], there exists a unique dominant weight μ such that $\mu = w\lambda$ for some w in W , and we have $\lambda \leq \mu$. Write $[m]_+ = [\lambda]_+ := \mu$ and call this dominant weight μ the *shape* of the monomial m or the weight λ .

For every monomial m of shape λ , Garsia and Stanton [14, proof of Theorem 9.4] showed that

$$m = \sum_{\substack{d \in \mathbb{Z}^r, w \in W: \\ \lambda_d + \lambda_{D(w)} = \lambda}} c_{d,w} a_1^{d_1} \cdots a_r^{d_r} z_w \quad (15)$$

is a linear combination of monomials whose shape is strictly less than λ , where $\lambda_d = d_1 \lambda_1 + \cdots + d_r \lambda_r$ and $c_{d,w} \in \mathbb{Z}$. It follows from induction that the descent monomials z_w form an $\mathbb{F}[\Lambda]^W$ -basis for $\mathbb{F}[\Lambda]$.

Lemma 5.1 *Suppose that γ is a simple root and λ is a weight such that $\langle \lambda, \gamma \rangle \geq 0$. If $0 \leq k \leq \langle \lambda, \gamma \rangle$ then $[\lambda - k\gamma]_+ \leq [\lambda]_+$, and the equality holds if and only if $k = 0$ or $\langle \lambda, \gamma \rangle$.*

Proof Let $\mu = \lambda - k\gamma$. If $k = 0$ or $\langle \lambda, \gamma \rangle$, then $\mu = \lambda$ or $s_\gamma \lambda$, and thus $[\mu]_+ = [\lambda]_+$ in either case.

Now assume $0 < k < \langle \lambda, \gamma \rangle$, and let $w\lambda$ and $u\mu$ be dominant for some w and u in W .

If $u\gamma > 0$ then $u\mu = uw^{-1}(w\lambda) - ku\gamma < uw^{-1}(w\lambda) \leq w\lambda$.

If $u\gamma < 0$ then

$$\begin{aligned} u\mu &= us_\gamma \cdot s_\gamma \lambda - ku\gamma \\ &= us_\gamma (\lambda - \langle \lambda, \gamma \rangle \gamma) - ku\gamma \\ &= us_\gamma \lambda + (\langle \lambda, \gamma \rangle - k)u\gamma \\ &< us_\gamma \lambda \leq w\lambda. \end{aligned}$$

Thus we are done. $\square$

Lemma 5.2 *Given a composition α of $r + 1$, let $\lambda_\alpha = \lambda_{D(\alpha)}$ and $z_\alpha = e^{\lambda_\alpha}$. If $w \in W$ has $D(w) \subseteq D(\alpha)$, then*

$$\bar{\pi}_w z_\alpha = e^{w\lambda_\alpha} + \sum_{[\lambda]_+ < \lambda_\alpha} c_\lambda e^\lambda, \quad c_\lambda \in \mathbb{Z}.$$

Moreover, $e^{w\lambda_\alpha}$ is a descent monomial if and only if $D(w) = D(\alpha)$.

Proof We prove the first assertion by induction on $\ell(w)$. If $\ell(w) = 0$ then we are done; otherwise $w = s_j u$ for some $j \in [r]$ and for some u with $\ell(u) < \ell(w)$. Since $D(u) \subseteq D(w) \subseteq D(\alpha)$, one has

$$\bar{\pi}_u z_\alpha = e^{u\lambda_\alpha} + \sum_{[\lambda]_+ < \lambda_\alpha} c_\lambda e^\lambda, \quad c_\lambda \in \mathbb{Z}.$$

Applying Lemma 5.1 to (14) (if the simple root γ_j satisfies $\langle \lambda, \gamma_j \rangle \leq 0$ then $\langle s_j \lambda, \gamma_j \rangle \geq 0$), one sees that

$$\bar{\pi}_j(e^\lambda) = \sum_{[\mu]_+ \leq [\lambda]_+} a_\mu e^\mu, \quad a_\mu \in \mathbb{Z}.$$

If we can show $\langle u\lambda_\alpha, \gamma_j \rangle > 0$, then applying Lemma 5.1 to the first case of (14) one has

$$\bar{\pi}_j e^{u\lambda_\alpha} = e^{s_j u\lambda_\alpha} + \sum_{[\mu]_+ < \lambda_\alpha} b_\mu e^\mu, \quad b_\mu \in \mathbb{Z}.$$

Combining these equations one obtains

$$\bar{\pi}_w z_\alpha = e^{w\lambda_\alpha} + \sum_{[\mu]_+ < \lambda_\alpha} b_\mu e^\mu + \sum_{[\lambda]_+ < \lambda_\alpha} c_\lambda \sum_{[\mu]_+ \leq [\lambda]_+} a_\mu e^\mu,$$

which gives the desired result.

Now we prove $\langle u\lambda_\alpha, \gamma_j \rangle > 0$. In fact, since $\ell(s_j u) > \ell(u)$, one has $u^{-1}(\gamma_j) > 0$, i.e.

$$u^{-1}(\gamma_j) = \sum_{i=1}^r m_i \gamma_i$$

for some nonnegative integers m_i . Applying $s_j u$ to both sides one gets

$$0 > -\gamma_j = \sum_{i=1}^r m_i s_j u(\gamma_i).$$

By the hypothesis $D(s_j u) \subseteq D(\alpha)$, if $i \notin D(\alpha)$ then $s_j u(\gamma_i) > 0$. This forces $m_i > 0$ for some $i \in D(\alpha)$, and thus

$$\langle u\lambda_\alpha, \gamma_j \rangle = \langle \lambda_\alpha, u^{-1}\gamma_j \rangle = \sum_{i \in D(\alpha)} m_i > 0.$$

Finally we consider when $e^{w\lambda_\alpha}$ is a descent monomial. If $D(w) = D(\alpha)$ then it is just the descent monomial of w . Conversely, if it is a descent monomial of some $u \in W$ then $w\lambda_\alpha = u\lambda_{D(u)}$. Since λ_α and $\lambda_{D(u)}$ are both in the fundamental Weyl chamber, the above equality implies that $\lambda_\alpha = \lambda_{D(u)}$ and $u^{-1}w$ is a product of simple reflections which all fix λ_α [19, Lemma 10.3B], i.e.

$$w = us_{j_1} \cdots s_{j_k}, \quad j_1, \dots, j_k \notin D(u) = D(\alpha).$$

Since $D(w) \subseteq D(\alpha) = D(u)$, none of $s_{j_1}, \dots, s_{j_k}$ is a descent of w , and thus it follows from the deletion property of W that w is a subword of some reduced expression of u , i.e. $w \leq u$ in Bruhat order. Similarly, it follows from $u = ws_{j_k} \cdots s_{j_1}$ that $u \leq w$ in Bruhat order. Thus $u = w$. $\square$

Theorem 5.3 *The coinvariant algebra $\mathbb{F}[\Lambda]/(a_1, \dots, a_r)$ has an $\mathbb{F}$ -basis $\{\pi_w e^{\lambda_{D(w)}} : w \in W\}$ and decomposes as*

$$\mathbb{F}[\Lambda]/(a_1, \dots, a_r) = \bigoplus_{\alpha \models r+1} H_W(0) \cdot \pi_{w_0(\alpha)z_\alpha}$$

where each direct summand $H_W(0) \cdot \pi_{w_0(\alpha)z_\alpha}$ has an $\mathbb{F}$ -basis

$$\{\pi_w z_\alpha : w \in [w_0(\alpha), w_1(\alpha)]\}$$

and is isomorphic to the projective indecomposable $H_W(0)$ -module $\mathbf{P}_\alpha$. Consequently, $\mathbb{F}[\Lambda]/(a_1, \dots, a_r)$ is isomorphic to the regular representation of $H_W(0)$.

Proof By Lemma 5.2, if one replaces the descent monomial z_w with the Demazure atom $\pi_w e^{\lambda_{D(w)}}$ in (15), the extra terms produced are of the form

$$c_{d,w} a_1^{d_1} \cdots a_r^{d_r} e^\mu$$

where $d = (d_1, \dots, d_r)$ and μ are weak compositions, and w is an element in W , satisfying $\lambda_d + \lambda_{D(w)} = \lambda$ and $[\mu] < \lambda_{D(w)}$. By the definition of $a_1, \dots, a_r$, one expands each term above as a linear combination of the monomials

$$e^{d_1 w_1 \lambda_1} \cdots e^{d_r w_r \lambda_r} e^\mu, \quad w_i \in W/W_{i^c}.$$

There exists $w \in W$ such that

$$\begin{aligned} [\mu + d_1 w_1 \lambda_1 + \cdots + d_r w_r \lambda_r] &= w(\mu + d_1 w_1 \lambda_1 + \cdots + d_r w_r \lambda_r) \\ &\leq [\mu] + d_1 \lambda_1 + \cdots + d_r \lambda_r \\ &< \lambda_{D(w)} + \lambda_d = \lambda. \end{aligned}$$

By induction on the shapes of the monomials, one shows that the Demazure atoms $\pi_w \lambda_{D(w)}$ for all w in W form an $\mathbb{F}[\Lambda]^W$ -basis for $\mathbb{F}[\Lambda]$, giving an $\mathbb{F}$ -basis for the coinvariant algebra $\mathbb{F}[\Lambda]/(a_1, \dots, a_r)$.

Let $\alpha \models r + 1$. For any u in W , using $\pi_i^2 = -\pi_i$ one shows by induction that $\pi_u \pi_{w_0(\alpha)} = \pi_w$ for some $w \geq w_0(\alpha)$ in the (left) weak order, which implies $D(w) \supseteq D(\alpha)$. On the other hand, if there exists $j \in D(w) \setminus D(\alpha)$, then $\pi_w z_\alpha = 0$ since $\pi_j z_\alpha = 0$ by (14). Hence $H_W(0) \cdot \pi_{w_0(\alpha)} z_\alpha$ has a basis $\{\pi_w z_\alpha : w \in [w_0(\alpha), w_1(\alpha)]\}$, and is isomorphic to $\mathbf{P}_\alpha$ via $\pi_w z_\alpha \mapsto T_w T'_{w_0(\alpha^c)}$ for all $w \in [w_0(\alpha), w_1(\alpha)]$. $\square$

Remark 5.4 Garsia and Stanton [14] pointed out a way to reduce the descent monomials in $\mathbb{F}[\Lambda]$ to the descent monomials in $\mathbb{F}[X]$ for type A . However, it does not give Theorem 4.5 directly from Theorem 5.3; instead, one should consider the Demazure operators on $\mathbb{F}[X(T)]$ where $X(T)$ is the character group of the subgroup T of diagonal matrices in $GL(n, \mathbb{F})$.

6 Flag varieties

In this section we assume $\mathbb{F}$ is a field of characteristic $p > 0$ and study the action of the 0-Hecke algebras on the (complete) flag varieties. Let G be a finite group of Lie type over a finite field $\mathbb{F}_q$ of characteristic p , with Borel subgroup B and Weyl group W . Assume that W is generated by simple reflections $s_1, \dots, s_r$. Every composition α of $r + 1$ corresponds to a *parabolic subgroup* $P_\alpha := BW_{D(\alpha)^c}B$ of G . The *partial flag variety* $1_{P_\alpha}^G$ is the induction of the trivial representation of P_α to G , or in other words, the $\mathbb{F}$ -span of all right P_α -cosets in G . Taking $\alpha = 1^{r+1}$ we have the *(complete) flag variety* 1_B^G .

For type A , one has $G = GL(n, \mathbb{F}_q)$, and if $\alpha = (\alpha_1, \dots, \alpha_\ell)$ is a composition of n , then P_α is the group of all block upper triangular matrices with invertible diagonal blocks of sizes $\alpha_1, \dots, \alpha_\ell$. Using the action of G on the vector space $V = \mathbb{F}^n$, one can identify $1_{P_\alpha}^G$ with the $\mathbb{F}$ -span of all partial flags of subspaces $0 \subset V_1 \subset \cdots \subset V_\ell = V$ satisfying $\dim V_i = \alpha_i$ for $i = 1, \dots, \ell$; in particular, 1_B^G is the $\mathbb{F}$ -span of all complete flags of V .

6.1 0-Hecke algebra action on 1_B^G

Given a subset $H \subseteq G$, let $\overline{H} = \sum_{h \in H} h$ be an element in the group algebra $\mathbb{Z}G$. Then $1_B^G = \overline{B} \cdot \mathbb{F}G$. By work of Kuhn [22], the endomorphism ring $\text{End}_{\mathbb{Z}G}(\overline{B} \cdot \mathbb{Z}G)$ has a basis $\{f_w : w \in W\}$, with f_w given by

$$f_w(\overline{B}) = \overline{BwB} = \overline{U_w w B}$$

where U_w is the product of the root subgroups of those positive roots which are sent to negative roots by w^{-1} (see e.g. [8, Proposition 1.7]). The endomorphism ring $\text{End}_{\mathbb{Z}G}(\bar{B} \cdot \mathbb{Z}G)$ is isomorphic to the Hecke algebra $H_W(q)$ of W with parameter $q = |U_{s_i}|$, since the relations satisfied by $\{f_w : w \in W\}$ are the same as those satisfied by the standard basis for $H_W(q)$. Working over a field $\mathbb{F}$ of characteristic p in which $q = 0$, we obtain a G -equivariant action of the 0-Hecke algebra $H_W(0)$ on 1_B^G by

$$T_w(\bar{B}g) := \overline{BwBg}, \quad \forall g \in G, \forall w \in W.$$

We will use left cosets in the next subsection, and in that case there is a similar right $H_W(0)$ -action.

Given a finite dimensional filtered $H_W(0)$ -module Q and a composition α of $r+1$, define Q_α to be the $\mathbb{F}$ -subspace of the elements in Q that are annihilated by T_j for all $j \notin D(\alpha)$, i.e.

$$Q_\alpha := \bigcap_{j \in D(\alpha)^c} \ker T_j.$$

The next lemma gives the simple composition factors of Q by inclusion–exclusion. We do not need any (nontrivial) filtration for Q in this section, but we will need it in the next section.

Lemma 6.1 *Given a finite dimensional filtered $H_W(0)$ -module Q and a composition α of $r+1$, the graded multiplicity of the simple $H_W(0)$ -module $\mathbf{C}_\alpha$ among the composition factors of Q is*

$$c_\alpha(Q) = \sum_{\beta \preceq \alpha} (-1)^{\ell(\alpha) - \ell(\beta)} \text{Hilb}(Q_\beta, t).$$

Proof Let $0 = Q_0 \subset Q_1 \subset \cdots \subset Q_k = Q$ be a composition series. We induct on the composition length k . The case $k = 0$ is trivial. Assume $k > 0$ below.

Suppose $Q/Q' \cong \mathbf{C}_\gamma$ for some $\gamma \models r+1$, i.e. there exists an element z in $Q \setminus Q'$ satisfying

$$T_i z \in \begin{cases} -z + Q', & \text{if } i \in D(\gamma), \\ Q', & \text{if } i \notin D(\gamma). \end{cases}$$

Let u be the longest element of the parabolic subgroup $W_{D(\gamma)^c}$ of W , and let

$$z' = T'_u z = \sum_{w \in W_{D(\gamma)^c}} T_w z.$$

Since $D(w) \subseteq D(\gamma)^c$ for all w in the above sum, we have $z' \in z + Q'$. Then any element in

$$Q = Q' \oplus \mathbb{F}z = Q' \oplus \mathbb{F}z'$$

can be written as $y + az'$ for some $y \in Q'$ and $a \in \mathbb{F}$. Since $D(u^{-1}) = D(\gamma)^c$, one has $T_i z' = T_i T'_u z = 0$ for all $i \notin D(\gamma)$. Consider an arbitrary composition β of $r+1$.

If $\gamma \preceq \beta$ then for any $i \notin D(\beta)$ we must have $i \notin D(\gamma)$ and thus $T_i(y + az') = T_i y$. It follows that $T_i(y + az') = 0$ if and only if $T_i y = 0$, i.e. $Q_\beta = Q'_\beta \oplus \mathbb{F}z'$.

If $\gamma \not\preceq \beta$ then there exists $i \in D(\gamma) \setminus D(\beta)$. Using $z' \in z + Q'$ we have

$$T_i(y + az') = T_i y + aT_i z' \in -az + Q'.$$

If $T_i(y + az') = 0$ then $a = 0$. This implies $Q_\beta = Q'_\beta$.

It follows that

$$\begin{aligned} c_\alpha(Q) &= \sum_{\beta \preceq \alpha} (-1)^{\ell(\alpha) - \ell(\beta)} \text{Hilb}(Q_\beta, t) \\ &= \sum_{\beta \preceq \alpha} (-1)^{\ell(\alpha) - \ell(\beta)} \text{Hilb}(Q'_\beta, t) + \sum_{\gamma \preceq \beta \preceq \alpha} (-1)^{\ell(\alpha) - \ell(\beta)} t^{\deg z'} \\ &= c_\alpha(Q') + \delta_{\alpha\gamma} \cdot t^{\deg z'}. \end{aligned}$$

On the other hand, by induction hypothesis, the graded multiplicity of $\mathbf{C}_\alpha$ in the composition factors of Q is also $c_\alpha(Q') + \delta_{\alpha\gamma} \cdot t^{\deg z'}$. Hence we are done. $\square$

Corollary 6.2 *If Q is a finite dimensional filtered $H_n(0)$ -module then*

$$\text{Ch}_t(Q) = \sum_{\alpha \models r+1} \text{Hilb}(Q_\alpha, t) M_\alpha.$$

Consequently, $\text{Ch}_t(Q) \in \text{QSym}[t]$ is a symmetric function, i.e. it lies in $\text{Sym}[t]$, if and only if

$$\text{Hilb}(Q_\alpha, t) = \text{Hilb}(Q_\beta, t) \quad \text{whenever } \beta \text{ is a rearrangement of } \alpha.$$

Proof Apply inclusion–exclusion to the previous lemma. $\square$

Remark 6.3 Lemma 6.1 and Corollary 6.2 hold for an arbitrary field $\mathbb{F}$.

Theorem 6.4 *The multiplicity of $\mathbf{C}_\alpha$ among the simple composition factors of 1_B^G is*

$$c_\alpha(1_B^G) = \sum_{w \in W: D(w^{-1}) = D(\alpha)} q^{\ell(w)}.$$

Proof Let $\overline{B}g$ be an element in 1_B^G where $g \in \mathbb{F}G$. If it is annihilated by T_j for all $j \in D(\alpha)^c$, then $\overline{B}w\overline{B}g = T_w(\overline{B}g) = 0$ for all w with $D(w) \cap D(\alpha)^c \neq \emptyset$, and in particular, for all w in $W_{D(\alpha)^c} \setminus \{1\}$. Hence

$$\overline{B}g = \overline{B}W_{D(\alpha)^c}\overline{B}g = \overline{P}_\alpha g \in 1_{P_\alpha}^G.$$

Conversely, $T_j(\overline{P}_\alpha g) = T_j T'_{w_0(D(\alpha)^c)}(\overline{B}g) = 0$ for all $j \in D(\alpha)^c$. Therefore $(1_B^G)_\alpha = 1_{P_\alpha}^G$. Applying Lemma 6.1 to 1_B^G (trivially filtered) gives

$$\begin{aligned} c_\alpha(1_B^G) &= \sum_{\beta \preccurlyeq \alpha} (-1)^{\ell(\alpha) - \ell(\beta)} |P_\alpha \setminus G| \\ &= \sum_{\beta \preccurlyeq \alpha} (-1)^{\ell(\alpha) - \ell(\beta)} \sum_{w \in W: D(w^{-1}) \subseteq D(\alpha)} |U_w| \\ &= \sum_{w \in W: D(w^{-1}) = D(\alpha)} q^{\ell(w)}. \end{aligned} \quad \square$$

Corollary 6.5 *If $G = GL(n, \mathbb{F}_q)$ then $\text{Ch}(1_B^G) = \tilde{H}_{1^n}(x; q)$.*

Proof For $G = GL(n, \mathbb{F}_q)$ we have $\ell(w) = \text{inv}(w)$ and thus equation (3) shows $c_\alpha(1_B^G) = r_\alpha(q)$. The result then follows from Corollary 4.9. $\square$

6.2 Decomposing the G -module 1_B^G by 0-Hecke algebra action

We consider the *homology representations* χ_q^α of G , which are the top homology of the *type-selected Tits buildings* of G , for all compositions $\alpha \models r + 1$. To give the explicit definitions, assume in this subsection that $1_{P_\alpha}^G$ is the $\mathbb{F}$ -span of the *left* P_α -cosets in G . Then 1_B^G admits a right $H_W(0)$ -action defined by $g\overline{B} \cdot T_w = g\overline{B}w\overline{B}$ for all $g \in G$ and $w \in W$. The left cosets gP_α for all $\alpha \models r + 1$ form a poset under reverse inclusion, giving an (abstract) simplicial complex called the *Tits building* and denoted by $\Delta = \Delta(G, B)$. The type of a face gP_α is $\tau(gP_\alpha) = D(\alpha)$, and every chamber gB has exactly one vertex of each type, i.e. $\Delta(G, B)$ is *balanced*. The chain complex of the *type-selected subcomplex*

$$\Delta_\alpha = \{F \in \Delta(G, B) : \tau(F) \subseteq D(\alpha)^c\}$$

gives rise to an exact sequence

$$0 \rightarrow \chi_q^\alpha \rightarrow 1_{P_\alpha}^G \xrightarrow{\partial} \bigoplus_{\beta \preccurlyeq_1 \alpha} 1_{P_\beta}^G \xrightarrow{\partial} \cdots \xrightarrow{\partial} 1_G^G \rightarrow 0 \quad (16)$$

where $\beta \preccurlyeq_1 \alpha$ means $\beta \models r + 1$ and $D(\beta) = D(\alpha) \setminus \{i\}$ for some $i \in D(\alpha)$. The boundary maps are given by

$$\partial : gP_\gamma \mapsto \sum_{\beta \preccurlyeq_1 \gamma} \pm gP_\beta$$

for all $\gamma \models r + 1$. The following decomposition of (left) G -modules is well-known (see e.g. Smith [29]):

$$1_B^G = \bigoplus_{\alpha \models r+1} \chi_q^\alpha. \quad (17)$$

On the other hand, Norton's decomposition of the 0-Hecke algebra $H_W(0)$ implies a decomposition of 1 into primitive orthogonal idempotents, i.e.

$$1 = \sum_{\alpha \models r+1} h_\alpha T_{w_0(\alpha)} T'_{w_0(\alpha^c)}, \quad h_\alpha \in H_W(0).$$

This decomposition of 1 into primitive orthogonal idempotents is explicitly given by Berg, Bergeron, Bhargava and Saliola [3], and is different from the one provided by Denton [7]. By the right action of $H_W(0)$ on 1_B^G , we have another decomposition of G -modules:

$$1_B^G = \bigoplus_{\alpha \models r+1} 1_B^G h_\alpha T_{w_0(\alpha)} T'_{w_0(\alpha^c)}. \quad (18)$$

Proposition 6.6 *The two G -module decompositions (17) and (18) are the same.*

Proof Comparing (17) with (18) one sees that it suffices to show $1_B^G h_\alpha T_{w_0(\alpha)} T'_{w_0(\alpha^c)} \subseteq \chi_q^\alpha$. Assume

$$\overline{B} h_\alpha T_{w_0(\alpha)} = \sum_i g_i \overline{B}, \quad g_i \in G.$$

For any $\beta \models r+1$ we have

$$\overline{B} T'_{w_0(\beta^c)} = \overline{B} W_{D(\beta)^c} \overline{B} = \overline{P}_\beta.$$

Hence

$$\overline{B} h_\alpha T_{w_0(\alpha)} T'_{w_0(\alpha^c)} = \sum_i g_i \overline{P}_\alpha \in 1_{P_\alpha}^G$$

and

$$\partial(\overline{B} h_\alpha T_{w_0(\alpha)} T'_{w_0(\alpha^c)}) = \sum_{\beta \preccurlyeq_1 \alpha} \pm \sum_i g_i \overline{P}_\beta = \sum_{\beta \preccurlyeq_1 \alpha} \pm \overline{B} h_\alpha T_{w_0(\alpha)} T'_{w_0(\beta^c)}.$$

If $\beta \preccurlyeq_1 \alpha$ then there exists $i \in D(\alpha) \cap D(\beta)^c$, and thus $T_{w_0(\alpha)} T'_{w_0(\beta^c)} = 0$. Therefore we are done. $\square$

One sees from (18) that χ_q^α is in general not a right $H_W(0)$ -submodule of 1_B^G . However, when $G = GL(n, \mathbb{F}_q)$, one sees that $\chi_q^{(n)}$ is the trivial representation and $\chi_q^{(1^n)}$ is the Steinberg representation of G [20], and both are right (isotypic) $H_W(0)$ -modules.

7 Coinvariant algebra of (G, B)

In this section we again assume $\mathbb{F}$ is a field of characteristic $p > 0$ and study the action of the 0-Hecke algebra $H_n(0)$ on the coinvariant algebra $\mathbb{F}[X]^B / (\mathbb{F}[X]_+^G)$ of

the pair (G, B) , where $G = GL(n, \mathbb{F}_q)$ with q being a power of p and B is the Borel subgroup of G .

Given a right $\mathbb{F}G$ -module M , there is an isomorphism

$$\begin{aligned} \text{Hom}_{\mathbb{F}G}(1_B^G, M) &\xrightarrow{\sim} M^B, \\ \phi &\mapsto \phi(\overline{B}) \end{aligned}$$

with inverse map given by $\phi_m(\overline{B}) = m$ for all $m \in M^B$. The left $H_n(0)$ -action $T_w \overline{B} = \overline{B} w B$ on 1_B^G commutes with the right G -action and induces a left action on $\text{Hom}_{\mathbb{F}G}(1_B^G, M)$ by

$$T_w(\phi)(\overline{B}) = \phi(T_{w^{-1}} \overline{B}) = \phi_m(\overline{B w^{-1} B}).$$

Hence we have a left $H_n(0)$ -action on M^B by

$$T_w(m) = T_w(\phi_m)(\overline{B}) = \phi_m(\overline{B w^{-1} B}) = \phi_m(\overline{B} w^{-1} \overline{U}_{w^{-1}}) = m w^{-1} \overline{U}_{w^{-1}}.$$

The group G has a left action on $\mathbb{F}[X]$ by linear substitution, and this can be turned into a right action by $f \cdot g = g^{-1} f$ for all $f \in \mathbb{F}[X]$ and $g \in G$. Thus $H_n(0)$ has a left action on $\mathbb{F}[X]^B$ by

$$T_w(f) = f \cdot w^{-1} \overline{U}_{w^{-1}} = \overline{U}_w w f, \quad \forall f \in \mathbb{F}[X]^B.$$

This action preserves the grading, and leaves the ideal $(\mathbb{F}[X]_+^G)$ invariant: if $h_i \in \mathbb{F}[X]_+^G$, $f_i \in \mathbb{F}[X]^B$, then

$$T_w\left(\sum_i h_i f_i\right) = \overline{U}_w w \left(\sum_i h_i f_i\right) = \sum_i h_i \overline{U}_w w(f_i).$$

Hence the coinvariant algebra $\mathbb{F}[X]^B / (\mathbb{F}[X]_+^G)$ of (G, B) becomes a graded $H_n(0)$ -module.

Lemma 7.1 *If $Q = \mathbb{F}[X]^B / (\mathbb{F}[X]_+^G)$ and α is a composition of n , then*

$$Q_\alpha := \bigcap_{j \in D(\alpha)^c} \ker T_j = \mathbb{F}[X]^{P_\alpha} / (\mathbb{F}[X]^G).$$

Proof If $f \in \mathbb{F}[X]^{P_\alpha}$, then for all $j \notin D(\alpha)$ one has $U_{s_j} s_j \subseteq P_\alpha$ and hence

$$T_j f = \overline{U}_{s_j} s_j f = |U_{s_j}| \cdot f = q f = 0.$$

Conversely, a B -invariant polynomial f gives rise to a P_α -invariant polynomial

$$\sum_{gB \in P_\alpha/B} g f = \sum_{w \in W_{D(\alpha)^c}} \overline{U}_w w f = T'_{w_0(D(\alpha)^c)} f.$$

If $T_j f$ belongs to the ideal $(\mathbb{F}[X]_+^G)$ for all $j \notin D(\alpha)$, so does $T_w f \in (\mathbb{F}[X]_+^G)$ for all $w \in W_{D(\alpha)^c} \setminus \{1\}$. Thus $T'_{w_0(D(\alpha)^c)} f - f \in (\mathbb{F}[X]_+^G)$ and we are done. $\square$

Theorem 7.2 *The graded quasi-symmetric characteristic of the $H_n(0)$ -module $\mathbb{F}[X]^B/(\mathbb{F}[X]_+^G)$ is*

$$\text{Ch}_t(\mathbb{F}[X]^B/(\mathbb{F}[X]_+^G)) = \sum_{\alpha \models n} \left[\begin{matrix} n \\ \alpha \end{matrix} \right]_{q,t} M_\alpha = \sum_{\alpha \models n} r_\alpha(q, t) F_\alpha.$$

Proof Let $Q = \mathbb{F}[X]^B/(\mathbb{F}[X]_+^G)$ and let $\alpha \models n$. It follows from Lemma 7.1 that

$$\text{Hilb}(Q_\alpha, t) = \text{Hilb}(\mathbb{F}[X]^{P_\alpha}/(\mathbb{F}[X]_+^G), t) = \left[\begin{matrix} n \\ \alpha \end{matrix} \right]_{q,t}.$$

Thus

$$c_\alpha(Q) = \sum_{\beta \preceq \alpha} (-1)^{\ell(\alpha) - \ell(\beta)} \left[\begin{matrix} n \\ \beta \end{matrix} \right]_{q,t} = r_\alpha(q, t).$$

Then the result follows immediately from Lemma 6.1. $\square$

8 Cohomology ring of Springer fibers

In Sect. 4 we showed that the coinvariant algebra of $\mathfrak{S}_n$ is an $H_n(0)$ -module whose graded quasi-symmetric characteristic is the modified Hall–Littlewood symmetric function indexed by the partition 1^n . Now we generalize this result to partitions of hook shapes.

Throughout this section a partition of n is denoted by $\mu = (\mu_1, \dots, \mu_n)$, where $0 \leq \mu_1 \leq \dots \leq \mu_n$. Denote $n(\mu) = \mu_{n-1} + 2\mu_{n-2} + \dots + (n-1)\mu_1$. One can view μ as a composition by dropping all the zero parts of μ . Then $n(\mu) = \text{maj}(\mu)$, where $\text{maj}(\alpha) = \sum_{i \in D(\alpha)} i$, for all compositions α .

Let V be an n -dimensional complex vector space. Fix a nilpotent matrix X_μ whose Jordan blocks are of size $\mu_1, \dots, \mu_\ell$. The *Springer fiber* $\mathcal{F}_\mu$ corresponding to the partition μ is the variety of all flags $0 \subset V_1 \subset \dots \subset V_n = V$ of subspaces $V_i \subseteq V$ satisfying $\dim V_i = i$ and $X_\mu(V_i) \subseteq V_{i-1}$. The cohomology ring of $\mathcal{F}_\mu$ is isomorphic to the ring $\mathbb{C}[X]/I_\mu$ for certain homogeneous ideal I_μ , and carries an $\mathfrak{S}_n$ -action that can be obtained from the $\mathfrak{S}_n$ -action on $\mathbb{C}[X]$. In particular, if $\mu = 1^n$ then $\mathcal{F}_\mu$ is the flag variety G/B and $\mathbb{C}[X]/I_\mu$ is the coinvariant algebra of $\mathfrak{S}_n$.

Theorem 8.1 (Hotta and Springer [18], Garsia and Procesi [13]) *The graded Frobenius characteristic of $\mathbb{C}[X]/I_\mu$ is the modified Hall–Littlewood symmetric function*

$$\tilde{H}_\mu(x; t) = \sum_{\lambda} t^{n(\mu)} K_{\lambda\mu}(t^{-1}) s_\lambda$$

where $K_{\lambda\mu}(t)$ is the Kostka–Foulkes polynomial.

To find an analogous result for the 0-Hecke algebras, let $R_\mu := \mathbb{F}[X]/I_\mu$ where $\mathbb{F}$ is an arbitrary field, and we consider the question of when the $H_n(0)$ -action on $\mathbb{F}[X]$

preserves the ideal I_μ . Recall the following construction of I_μ by Tanisaki [32]. Let the conjugate of a partition μ of n be $\mu' = (0 \leq \mu'_1 \leq \dots \leq \mu'_n)$. Note that the height of the Ferrers diagram of μ is $h = h(\mu) := \mu'_n$. Let

$$d_k(\mu) = \mu'_1 + \dots + \mu'_k, \quad k = 1, \dots, n.$$

Then the ideal I_μ is generated by

$$\{e_r(S) : k \geq r > k - d_k(\mu), |S| = k, S \subseteq \{x_1, \dots, x_n\}\} \quad (19)$$

where $e_r(S)$ is the r th elementary symmetric function in the set S of variables. See also Garsia and Procesi [13].

Proposition 8.2 *The Demazure operators preserve the ideal I_μ if and only if μ is a hook.*

Proof First assume $\mu = (0^{n-h}, 1^{h-1}, n-h+1)$ is a hook. Then $\mu' = (0^{h-1}, 1^{n-h}, h)$ and so

$$(1 - d_1(\mu), 2 - d_2(\mu), \dots, n - d_n(\mu)) = (1, 2, \dots, h-1, h-1, \dots, h-1, 0).$$

It follows that the ideal I_μ is generated by the elementary symmetric functions $e_1, \dots, e_n$, together with the following partial elementary symmetric functions:

$$\{e_r(S) : r = h, \dots, k, S \subseteq \{x_1, \dots, x_n\}, |S| = k, k = h, \dots, n-1\}.$$

This implies that I_μ can be generated by $e_1, \dots, e_n$ and all monomials in

$$\mathcal{M}_h = \{x_{i_1} \cdots x_{i_h} : 0 \leq i_1 < \dots < i_h \leq n\}.$$

By (10), if $x_{i_1} \cdots x_{i_h}$ is in $\mathcal{M}_h$ and f is an arbitrary monomial, then $\pi_i(x_{i_1} \cdots x_{i_h} f)$ is divisible by some element in $\mathcal{M}_h$. Thus the ideal I_μ is preserved by the Demazure operators.

Now assume μ is not a hook. Then $\mu'_{n-1} \geq 2$ and thus

$$\begin{aligned} k - d_k(\mu) &= k - n + n - d_k(\mu) \\ &= k - n + \mu'_n + \mu'_{n-1} + \dots + \mu'_{k+1} \\ &\geq k - n + \mu'_n + 2 + 1 + \dots + 1 \\ &= \mu'_n = h \end{aligned}$$

for $k = n-2, \dots, n - \mu_1 + 1$. One also sees that

$$k - d_k(\mu) = \begin{cases} 0, & k = n, \\ h-1, & k = n-1, \\ k, & n - \mu_1 \geq k \geq 1. \end{cases}$$

Thus the only elements in the generating set (19) that have degree no more than h are $e_1, \dots, e_h$ and those $e_h(S)$ with $|S| = n-1$.

Suppose to the contrary that I_μ is preserved by Demazure operators. Since

$$e_h(x_1, \dots, x_{n-1}) = x_{n-1}e_{h-1}(x_1, \dots, x_{n-2}) + e_h(x_1, \dots, x_{n-2}) \in I_\mu$$

we have

$$\overline{\pi}_{n-1}e_h(x_1, \dots, x_{n-1}) = x_n e_{h-1}(x_1, \dots, x_{n-2}) \in I_\mu$$

and thus

$$e_h(x_1, \dots, x_{n-2}, x_n) - \overline{\pi}_{n-1}e_h(x_1, \dots, x_{n-1}) = e_h(x_1, \dots, x_{n-2}) \in I_\mu.$$

Repeating this process one obtains $e_h(x_1, \dots, x_h) = x_1 \cdots x_h \in I_\mu$. Then applying the Demazure operators to $x_1 \cdots x_h$ gives $x_{i_1} \cdots x_{i_h} \in I_\mu$ whenever $1 \leq i_1 < \cdots < i_h \leq n$. Considering the degree we have

$$x_{i_1} \cdots x_{i_h} = \sum_{i=1}^h f_i e_i + \sum_{|S|=n-1} c_S e_h(S)$$

where $f_i \in \mathbb{F}[X]$ is homogeneous of degree $h-i$ and $c_S \in \mathbb{F}$. It follows that $U \subseteq U'$, where U and U' are the $\mathbb{F}$ -subspaces of the coinvariant algebra $\mathbb{F}[X]/(e_1, \dots, e_n)$ spanned by

$$\{x_{i_1} \cdots x_{i_h} : 1 \leq i_1 < \cdots < i_h \leq n-1\} \quad \text{and} \quad \{e_h(S) : |S| = n-1\}.$$

It is well known that all divisors of $x_1^{n-1} x_2^{n-2} \cdots x_{n-1}$ form a basis for $\mathbb{F}[X]/(e_1, \dots, e_n)$, i.e. the *Artin basis*. Thus

$$n = \binom{n}{n-1} \geq \dim U' \geq \dim U = \binom{n-1}{h} \geq \binom{n-1}{2}$$

where the last inequality follows from $2 \leq h \leq n-2$ since μ is a hook. Therefore we must have $n = 4$ and $h = 2$. In that case it is easy to check $x_1 x_2 \notin I_{(2,2)}$. Hence we are done. $\square$

Theorem 8.3 Assume $\mu = (0^{n-h}, 1^{h-1}, n-h+1)$ is a hook and view it as a composition by removing all zeros. Then the $H_n(0)$ -module R_μ is a direct sum of the projective indecomposable $H_n(0)$ -modules $\mathbf{P}_\alpha$ for all compositions $\alpha \preceq \mu$, i.e.

$$R_\mu \cong \bigoplus_{\alpha \preceq \mu} \mathbf{P}_\alpha.$$

Proof By the proof of the previous proposition, I_μ is generated by $e_1, \dots, e_n$ and

$$\mathcal{M}_h = \{x_{i_1} \cdots x_{i_h} : 1 \leq i_1 < \cdots < i_h \leq n\}.$$

Thus R_μ is the quotient of $\mathbb{F}[X]/(\mathbb{F}[X]_+^{\mathfrak{S}_n})$ by its ideal generated by $\mathcal{M}_h$. By Theorem 4.5, it suffices to show that $\mathbb{F}[X]/I_\mu$ has a basis given by

$$\{\overline{\pi}_{wX_{D(w)}} : w \in \mathfrak{S}_n, D(w) \subseteq D(\mu)\}. \quad (20)$$

If $D(w) \not\subseteq D(\mu) = \{1, 2, \dots, h-1\}$, i.e. w has a descent $i \geq h$, then $x_{D(w)}$ contains at least h distinct variables, and so do all monomials in $\overline{\pi}_w x_{D(w)}$ by (10). Thus $\mathbb{F}[X]/I_\mu$ is spanned by (20).

To show (20) is linearly independent, assume

$$\sum_{D(w) \subseteq D(\mu)} c_w \overline{\pi}_w x_{D(w)} \in I_\mu, \quad c_w \in \mathbb{F}.$$

For any polynomial $f \in \mathbb{F}[X]$, let $[f]_h$ be the polynomial obtained from f by removing all terms divisible by some element in $\mathcal{M}_h$. It follows that

$$\sum_{D(w) \subseteq D(\mu)} c_w [\overline{\pi}_w x_{D(w)}]_h \in (e_1, \dots, e_{h-1}).$$

If $D(w) \subseteq D(\mu) = [h-1]$ then the leading term of $[\overline{\pi}_w x_{D(w)}]_h$ under “ $<$ ” is still the descent monomial $w x_{D(w)}$. By Lemma 4.4,

$$\{[\overline{\pi}_w x_{D(w)}]_h : w \in \mathfrak{S}_n, D(w) \subseteq D(\mu)\}$$

gives a linearly independent set in $\mathbb{F}[X]/(\mathbb{F}[X]_+^{\mathfrak{S}_n})$. This forces $c_w = 0$ whenever $D(w) \not\subseteq D(\mu)$. $\square$

By work of Bergeron and Zabrocki [4], the modified Hall–Littlewood functions $\tilde{H}_\mu(x; t)$ have the following non-commutative analogue lying in $\mathbf{NSym}[t]$ for all compositions α :

$$\tilde{\mathbf{H}}_\alpha(\mathbf{x}; t) := \sum_{\beta \preccurlyeq \alpha} t^{\text{maj}(\beta)} \mathbf{s}_\beta.$$

Corollary 8.4 *Assume μ is a hook. Then*

$$\mathbf{ch}_t(R_\mu) = \sum_{\alpha \preccurlyeq \mu} t^{\text{maj}(\alpha)} \mathbf{s}_\alpha = \tilde{\mathbf{H}}_\mu(X; t), \quad (21)$$

$$\text{Ch}_t(R_\mu) = \sum_{\alpha \preccurlyeq \mu} t^{\text{maj}(\alpha)} s_\alpha = \tilde{H}_\mu(x; t). \quad (22)$$

Proof Theorem 8.3 immediately implies the graded non-commutative characteristic of R_μ , whose commutative image is the graded quasi-symmetric characteristic of R_μ . One can check that $\text{Ch}_t(R_\mu) = \tilde{H}_\mu(x; t)$ when μ is a hook, by using Haglund’s combinatorial formula [15] for the modified Macdonald polynomials $\tilde{H}_\mu(x; q, t)$ and taking $q \rightarrow 0$. $\square$

Remark 8.5

- (i) We recover the results in Sect. 4 by taking $\mu = 1^n$.
- (ii) Using certain difference operators, Hivert [16] defined a non-commutative analogue of the Hall–Littlewood functions, which is in general different from the non-commutative analogue of Bergeron and Zabrocki. However, they are the same when μ is a hook!

- (iii) It is not clear to the author why the results are nice only in the hook case, except for a naive explanation: the hooks are the only diagrams that belong to both the family of the Ferrers diagrams of partitions and the family of ribbon diagrams of compositions.

9 Questions for future research

9.1 Equidistribution of the inversion number and major index

The equidistribution of inv and maj was first proved on permutations of multisets by P.A. MacMahon in the 1910s; applying an inclusion–exclusion would give their equidistribution on inverse descent classes of $\mathfrak{S}_n$. However, the first proof for the latter result appearing in the literature was by Foata and Schützenberger [11] in 1970, using a bijection constructed earlier by Foata [10]. Is there an algebraic proof from the (q, t) -bigraded characteristic of $\mathbb{F}[X]/(\mathbb{F}[X]_+^{\mathfrak{S}_n})$, which is given in Corollary 4.9(i) and involves inv , maj , and inverse descents?

9.2 Decompositions of 1_B^G and $\mathbb{F}[X]^B/(\mathbb{F}[X]_+^G)$

In Sect. 6 we studied an $H_W(0)$ -action on the flag variety 1_B^G and found its simple composition factors, but we do *not* know the decomposition of 1_B^G into indecomposable $H_W(0)$ -modules. Assume $G = GL(n, \mathbb{F}_q)$ below. Computations show that 1_B^G is in general *not* projective, although its quasi-symmetric characteristic is always symmetric.

The coinvariant algebra $\mathbb{F}[X]^B/(\mathbb{F}[X]_+^G)$ is *not* a projective $H_n(0)$ -module either, since its graded quasi-symmetric characteristic is *not* even symmetric (see the definition of the (q, t) -multinomial coefficients). To find its decomposition, it will be helpful to know more (nonprojective) indecomposable $H_n(0)$ -modules (there are infinitely many, and some were studied by Duchamp, Hivert, and Thibon [9]).

Another question is to find a q -analogue of the Demazure operators, which might give another $H_n(0)$ -action on $\mathbb{F}[X]^B/(\mathbb{F}[X]_+^G)$.

9.3 Coincidence of Frobenius type characteristics

For $G = GL(n, \mathbb{C})$, the complex flag variety 1_B^G has its cohomology ring isomorphic to the coinvariant algebra of $\mathfrak{S}_n$, whose graded Frobenius characteristic and graded quasi-symmetric characteristic both equal the modified Hall–Littlewood symmetric function $\tilde{H}_{1^n}(x; t)$. For $G = GL(n, \mathbb{F}_q)$, the flag variety 1_B^G itself, when defined over a field of characteristic $p \mid q$, is also an $H_n(0)$ -module whose quasi-symmetric characteristic equals $\tilde{H}_{1^n}(x; q)$. Is there a better explanation for the coincidence of these Frobenius type characteristics?

Acknowledgements The author is grateful to Victor Reiner for asking questions which led to the research, for giving many helpful suggestions, and for his support from NSF grant DMS-1001933. He is indebted to Andrew Berget, Tom Denton, Anne Schilling, and Monica Vazirani for interesting conversations and hospitality during the visit to UC Davis. He thanks Marcelo Aguiar, Sami Assaf, Nantel Bergeron, Adriano Garsia, Patricia Hersh, Liping Li, Sarah Mason, Dennis Stanton and Peter Webb for their helpful suggestions and comments.

References

- Allen, E.E.: The descent monomials and a basis for the diagonally symmetric polynomials. *J. Algebr. Comb.* **3**, 5–16 (1994)
- Assem, I., Simson, D., Skowroński, A.: Elements of the Representation Theory of Associative Algebras, vol. 1: Techniques of Representation Theory. London Mathematical Society Student Texts, vol. 65. Cambridge University Press, Cambridge (2006)
- Berg, C., Bergeron, N., Bhargava, S., Saliola, F.: Primitive orthogonal idempotents for R -trivial monoids. *J. Algebra* **348**, 446–461 (2011)
- Bergeron, N., Zabrocki, M.: q and q, t -analogues of non-commutative symmetric functions. *Discrete Math.* **298**, 79–103 (2005)
- Björner, A., Wachs, M.: Generalized quotients in Coxeter groups. *Trans. Am. Math. Soc.* **308**, 1–37 (1988)
- Demazure, M.: Désingularisation des variétés de Schubert généralisées. *Ann. Sci. Éc. Norm. Supér.* **7**, 53–88 (1974)
- Denton, T.: A combinatorial formula for orthogonal idempotents in the 0-Hecke algebra of the symmetric group. *Electron. J. Comb.* **18** (2011)
- Digne, F., Michel, J.: Representations of Finite Groups of Lie Type. London Math. Soc. Student Texts, vol. 21. Cambridge University Press, Cambridge (1991)
- Duchamp, G., Hivert, F., Thibon, J.-Y.: Noncommutative symmetric functions VI: Free quasi-symmetric functions and related algebras. *Int. J. Algebra Comput.* **12**, 671–717 (2002)
- Foata, D.: On the Netto inversion number of a sequence. *Proc. Am. Math. Soc.* **19**, 236–240 (1968)
- Foata, D., Schützenberger, M.-P.: Major index and inversion number of permutations. *Math. Nachr.* **83**, 143–159 (1970)
- Garsia, A.: Combinatorial methods in the theory of Cohen–Macaulay rings. *Adv. Math.* **38**, 229–266 (1980)
- Garsia, A., Procesi, C.: On certain graded S_n -modules and the q -Kostka polynomials. *Adv. Math.* **94**, 82–138 (1992)
- Garsia, A., Stanton, D.: Group actions of Stanley–Reisner rings and invariant of permutation groups. *Adv. Math.* **51**, 107–201 (1984)
- Haglund, J., Haiman, M., Loehr, N.: A combinatorial formula for Macdonald polynomials. *J. Am. Math. Soc.* **18**, 735–761 (2005)
- Hivert, F.: Hecke algebras, difference operators, and quasi-symmetric functions. *Adv. Math.* **155**, 181–238 (2000)
- Hivert, F., Novelli, J.-C., Thibon, J.-Y.: Yang–Baxter bases of 0-Hecke algebras and representation theory of 0-Ariki–Koike–Shoji algebras. *Adv. Math.* **205**, 504–548 (2006)
- Hotta, R., Springer, T.A.: A specialization theorem for certain Weyl group representations and an application to the Green polynomials of unitary groups. *Invent. Math.* **41**, 113–127 (1977)
- Humphreys, J.: Introduction to Lie Algebras and Representation Theory. Graduate Texts in Mathematics, vol. 9. Springer, New York (1978)
- Khammash, A.A.: On the homological construction of the Steinberg representation. *J. Pure Appl. Algebra* **87**, 17–21 (1993)
- Krob, D., Thibon, J.-Y.: Noncommutative symmetric functions IV: Quantum linear groups and Hecke algebras at $q = 0$. *J. Algebr. Comb.* **6**, 339–376 (1997)
- Kuhn, N.: The modular Hecke algebra and Steinberg representation of finite Chevalley groups. *J. Algebra* **91**, 125–141 (1984)
- Kumar, S.: Demazure character formula in arbitrary Kac–Moody setting. *Invent. Math.* **89**, 395–423 (1987)
- Lascoux, A., Schützenberger, M.-P.: Sur une conjecture de H. O. Foulkes. *C. R. Acad. Sci. Paris Sér. A-B* **286**(7), A323–A324 (1978)
- Mason, S.: An explicit construction of type A Demazure atoms. *J. Algebr. Comb.* **29**, 295–313 (2009)
- Norton, P.N.: 0-Hecke algebras. *J. Aust. Math. Soc. A* **27**, 337–357 (1979)
- Reiner, V., Stanton, D.: (q, t) -analogues and $GL_n(\mathbb{F}_q)$. *J. Algebr. Comb.* **31**, 411–454 (2010)
- Schützenberger, M.-P.: La correspondance de Robinson. *Lecture Notes in Math.*, vol. 579, pp. 59–113. Springer, Berlin (1977)
- Smith, S.D.: On decomposition of modular representations from Cohen–Macaulay geometries. *J. Algebra* **131**, 598–625 (1990)

30. Stanley, R.: Ordered Structures and Partitions. Mem. Am. Math. Soc., vol. 119 (1972)
31. Steinberg, R.: On a theorem of Pittie. Topology **14**, 173–177 (1975)
32. Tanisaki, T.: Defining ideals of the closures of conjugacy classes and representations of the Weyl groups. Tohoku Math. J. **34**, 575–585 (1982)