Erratum to: Nitrogen detected TROSY at high field yields high resolution and sensitivity for protein NMR

Koh Takeuchi^{1,2} · Haribabu Arthanari³ · Ichio Shimada^{1,4} · Gerhard Wagner³

Published online: 11 December 2017
© Springer Science+Business Media B.V. 2017

Erratum to: J Biomol NMR (2015) 63(4):323–331
<https://doi.org/10.1007/s10858-015-9991-y>

The authors regret, a mistake appeared in the supplement of this paper. The fourth term of the Eq. 1 in the supplemental material reads $6J(\omega_S)$, instead of $6J(\omega_I)$. Furthermore, the lower left panel for the supplemental Figure S2 was not for a protein with a rotational correlation time (τ_c) of 7 ns but

for a protein with a τ_c of 20 ns. The corrected figure corresponding to a τ_c of a 7 ns protein is in the supplementary material. These changes do not influence the conclusions of the paper. The authors apologize for any inconvenience caused. The correct version of the Electronic supplementary material is given in the link below.

The online version of the original article can be found under
<https://doi.org/10.1007/s10858-015-9991-y>.

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s10858-017-0131-8>) contains supplementary material, which is available to authorized users.

-
- ✉ Ichio Shimada
shimada@iw-nmr.f.u-tokyo.ac.jp
- ✉ Gerhard Wagner
gerhard_wagner@hms.harvard.edu

- ¹ Molecular Profiling Research Center for Drug Discovery, National Institute for Advanced Industrial Science and Technology, Tokyo 135-0063, Japan
- ² PRESTO, JST, Tokyo 135-0063, Japan
- ³ Department of Biochemistry and Molecular Pharmacology, Harvard Medical School, 240 Longwood Avenue, Boston, MA 02115, USA
- ⁴ Graduate Schools of Pharmaceutical Sciences, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan