ERRATUM

Erratum to: Defective Leukocyte Adhesion and Chemotaxis Contributes to Combined Immunodeficiency in Humans with Autosomal Recessive MST1 Deficiency

Tarana Singh Dang¹ • Joseph D. P. Willet¹ • Helen R. Griffin² • Neil V. Morgan³ • Graeme O'Boyle¹ • Peter D. Arkwright⁴ • Stephen M. Hughes⁴ • Mario Abinun^{1,5} • Louise J. Tee³ • Dawn Barge⁶ • Karin R. Engelhardt¹ • Michael Jackson⁵ • Andrew J. Cant^{1,5} • Eamonn R. Maher^{3,7} • Mauro Santibanez Koref² • Louise N. Reynard¹ • Simi Ali¹ • Sophie Hambleton^{1,5}

Published online: 3 March 2016
© Springer Science+Business Media New York 2016

Erratum to: J Clin Immunol (2016)
DOI 10.1007/s10875-016-0232-2

The original version of this article unfortunately contained mistakes in Table 1 and author's contribution. The correct version is presented on the next page.

Tarana Singh Dang and Joseph D. P. Willet contributed equally to this work.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10875-016-0232-2>.

✉ Sophie Hambleton
sophie.hambleton@newcastle.ac.uk

¹ Institute for Cellular Medicine, Newcastle University, Newcastle upon Tyne NE2 4HH, UK

² Institute of Genetic Medicine, Newcastle University, Newcastle upon Tyne, UK

³ School of Clinical and Experimental Medicine, University of Birmingham, Birmingham, UK

⁴ Department of Paediatric Allergy & Immunology, Royal Manchester Children's Hospital, University of Manchester, Manchester, UK

⁵ Great North Children's Hospital, Newcastle upon Tyne Hospitals NHS Foundation Trust, Newcastle upon Tyne, UK

⁶ Blood Sciences Flow Cytometry Laboratory, Newcastle upon Tyne Hospitals NHS Foundation Trust, Newcastle upon Tyne, UK

⁷ Department of Medical Genetics, University of Cambridge, Cambridge, UK

Table 1 Clinical features and laboratory values for patients, demonstrating profound CD4-lymphopenia and reduced lymphoproliferative responses

Feature ^a	P1	P2	P3	Normal range ^c
Clinical features				
Eczema & skin infections	+	+	–	
Recurrent sinopulmonary infection	+	+	–	
Oral candidiasis	–	+	–	
Cryptosporidiosis	+	+	+	
Viral infections:				
<i>Molluscum contagiosum</i>	+	+	+	
Recurrent VZV	–	–	+	
Severe 1° HSV	+	+	–	
Chronic EBV viremia	+	+	+	
EBV LPD	+	+	–	
Flow cytometric immunophenotyping (<i>absolute cell counts per µl except where % indicated</i>)				
CD3 ⁺ T cells	815	850	921	1400–8000
CD4 ⁺ T cells	200	268	368	900–5500
^b naïve CD4 ⁺ T cells	0	17	18	
CD8 ⁺ T cells	512	592	523	400–2300
^b naïve CD8 ⁺ T cells	16	17	28	
TCR-γδ ⁺ (% of T cells)	28	7	15	
HLA-DR ⁺ (% of T cells)	84	77	62	
CD16 ⁺ /CD56 ⁺ NK cells	192	147	298	100–1400
CD19 ⁺ B cells	439	897	1355	600–3100
CD27 ⁺ IgD [–] (% of B cells)	2	3	5	
Lymphocyte proliferation assay				
PHA patient (cpm)	11,326	34,343	67,254	
PHA control (cpm)	148,110	122,215	101,164	
SI patient	82	29	94	
SI control	445	407	349	
Immunoglobulins				
<i>IgG</i>	18.3	28.8	17.8	3.0–13.3 g/L
<i>IgA</i>	2.2	4.47	0.7	0.3–1.29 g/L
<i>IgM</i>	0.6	1.13	0.9	0.43–1.9 g/L
<i>Pneumococcal ab</i>	<3	17	88	20–200 mg/L
<i>Tetanus ab</i>	0.88	1.55	0.49	0.1–10 IU/ml
<i>Hemophilus B ab</i>	NT	>9.0	NT	1.0–20 mg/L

VZV Varicella zoster virus, HSV Herpes simplex virus, EBV Epstein Barr Virus, LPD lymphoproliferative disease, PHA phytohemagglutinin, cpm counts per minute, SI stimulation index, ab specific antibody, NT not tested

^a Laboratory evaluation was performed at 5.5 years (P1), 3.4 years (P2), 1.7 years (P3)

^b Naïve T cells were defined as CD3 + CD27 + CD45RA⁺ and as either CD4⁺ or CD4[–]

^c Normal ranges are age matched to P3