Corrigendum to: Why is the relative abundance of *Asterionella formosa* increasing in a Boreal Shield lake as nutrient levels decline?

Branaavan Sivarajah · Kathleen M. Rühland · Andrew L. Labaj ·
Andrew M. Paterson · John P. Smol

Published online: 17 May 2016
© Springer Science+Business Media Dordrecht 2016

**Corrigendum to: J Paleolimnol (2016) 55:357–367
DOI 10.1007/s10933-016-9886-2**

Upon reading the published version of Sivarajah et al. (2016), we noted the last two sentences in the Results (p. 361, right-hand column) should read:

The increase in the relative abundance of *A. formosa* shows no relationship to the decline in

TIN (Spearman Rank correlation: $r_s = -0.26$, $N = 14$, $p = 0.37$) and is marginally related (but not statistically significant at $\alpha = 0.05$) to the increase in mean annual air temperature (Spearman Rank correlation: $r_s = 0.40$, $N = 19$, $p = 0.09$).

B. Sivarajah · K. M. Rühland · A. L. Labaj ·
J. P. Smol (✉)
Paleoecological Environmental Assessment and Research
Laboratory, Department of Biology, Queen's University,
Kingston, ON, Canada
e-mail: smolj@queensu.ca

A. M. Paterson
Dorset Environmental Science Centre, Ontario Ministry
of the Environment and Climate Change, Dorset, ON,
Canada