Erratum to: New Pb-based Superconductor with the 1222 Structure in the $(\text{Pb}_{0.75}\text{P}_{0.25})\text{Sr}_2(\text{Eu}_{1.9-x}\text{Ce}_x\text{Sr}_{0.1})\text{Cu}_2\text{O}_z$ System

H. Sasakura · Y. Akagi · M. Tanaka · S. Tsukui ·
M. Adachi

Published online: 10 August 2011
© Springer Science+Business Media, LLC 2011

Erratum to: J Supercond Nov Magn (2011) 24:1623–1626

DOI [10.1007/s10948-010-1068-2](https://doi.org/10.1007/s10948-010-1068-2)

Two errors occur on p. 1624 in *Journal of Superconductivity and Novel Magnetism*, Vol. 24, No. 5, pp. 1623–1626 (2011). In Fig. 1, the indices of the plane (10 $\overline{10}$) and (114) should be corrected to (00 $\overline{10}$) and (109), respectively. Here is the correct Fig. 1.

Fig. 1 X-ray diffraction patterns for the sample with the composition of $0.3 \leq x \leq 0.7$ in the present $(\text{Pb}_{0.75}\text{P}_{0.25})\text{Sr}_2(\text{Eu}_{1.9-x}\text{Ce}_x\text{Sr}_{0.1})\text{Cu}_2\text{O}_z$ system. For an example, diffraction peaks for the sample with $x = 0.7$ are indexed on the basis of a tetragonal unit cell with the lattice parameters of $a = 0.3851$ nm and $c = 2.922$ nm. The indexed peaks are attributable to those of the Pb-1222 phase and each value of the lattice parameters is almost the same as that of the Pb-1222 phase. But the sample with $x = 0.2$ is found to deviate from Pb-1222 single phase

The online version of the original article can be found under doi:[10.1007/s10948-010-1068-2](https://doi.org/10.1007/s10948-010-1068-2).

H. Sasakura (✉) · Y. Akagi
Department of Physics, Hamamatsu University School
of Medicine, 1-20-1 Handayama, Higashi-ku, Hamamatsu-shi,
Shizuoka 431-3192, Japan
e-mail: sasakura@hama-med.ac.jp

M. Tanaka
Department of Physics, Shizuoka University, 836 Ohya,
Suruga-ku, Shizuoka 422-8529, Japan

S. Tsukui · M. Adachi
Department of Chemical Engineering, Graduate School
of Engineering, Osaka Prefecture University, Naka-ku, Sakai-shi,
Osaka 599-8531, Japan