Erratum to: Attenuation coefficients of Rayleigh and L_g waves

Igor B. Morozov

Published online: 15 September 2010
© Springer Science+Business Media B.V. 2010

Erratum to: J Seismol (2010) DOI 10.1007/s10950-010-9196-5

In Morozov (2010), there was an error in plotting the long-period attenuation coefficients in Fig. 3b. The correct Fig. 3 is given here. Also, the intercept value of the interpreted trend (dashed line

in Fig. 3b) stated in the caption and in the text should be $-8 \cdot 10^{-5} \text{ s}^{-1}$. These corrections do not affect any conclusions of the paper. In fact, in the corrected plot, the suggested piecewise-linear pattern of attenuation coefficients becomes more pronounced even for normal modes.

Fig. 3 Spheroidal-mode (black dots) and Rayleigh-wave (other symbols) attenuation data from IGPP reference model web site: **a** in the original $1,000 Q^{-1}$ form; **b** transformed to $\chi(f)$. Dashed line shows the interpreted linear

trend $\chi(f) = -8 \cdot 10^{-5} + \pi f/Q_e \text{ [s}^{-1}\text{]}$, with $Q_e = 84$. f_c is the “cross-over” frequency at which the change from surface-wave to normal-mode regime occurs

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10950-010-9196-5>.

I. B. Morozov (✉)
Department of Geological Sciences,
University of Saskatchewan, Saskatoon,
SK S7N 5E2, Canada
e-mail: igor.morozov@usask.ca

Reference

Morozov IB (2010) Attenuation coefficients of Rayleigh and L_g waves. J Seismol. doi:10.1007/s10950-010-9196-5