ERRATUM

To the article “Measure Convolution Semigroups and Noninfinitely Divisible Probability Distributions” by Aubrey Wulfsohn (*Journal of Mathematical Sciences*, Vol. 131, No. 3, pp. 5682–5696, December, 2005).

The Introduction to this article, which was inadvertently omitted, appears below:

Introduction

Let μ be a probability measure (or corresponding random variable) such that all moments exist. Knowledge of the moments μ_n is not sufficient to determine infinite divisibility of μ ; we show also that infinitely divisible and, in particular, lognormal distributions lose infinite divisibility when censored in certain ways even if all moments are arbitrarily close to those of the uncensored distribution. The moments of a composition of k copies of μ are expressed as combinatorial compositions of the μ_n . We express the moments of the compositions in the context of occupancy problems, arranging n balls in k cells; the classical convolution is described by Maxwell–Boltzmann statistics and is multinomial. For certain noninfinitely divisible measures with moments increasing fast enough the indexing of a k -cell combinatorial composition is extended to indexing by nonnegative real t , and we construct classical convolution measure semigroups from amongst the t -indexed classes. We prove also that when a random variable with infinitely divisible distribution is embedded in a Lévy process (Y_t) , then the t -indexed Maxwell–Boltzmann is the law of Y_t . In order to get moment-based multinomial compositions indexed by a continuum, we use random measures and random distributions rather than random variables. An alternative approach to embedability of a noninfinitely divisible μ is by considering nonclassical convolution measure semigroups; for example, embedding μ in a Boolean convolution measure semigroup and retaining the multinomial character of the moments. Embedding μ in an Urbanik generalized convolution measure semigroup one loses the multinomial character of the moments.