FOREWORD

V. M. Manuilov and A. S. Mishchenko

UDC 51.0

This issue and subsequent ones present talks given at the international conference “Topology, analysis and applications to mathematical physics” dedicated to the memory of Professor Yu. P. Solovyov held in Moscow on February 14–19, 2005. The conference gathered many topologists from Russia, from the republics of the former Soviet Union, and from other countries, who were students of Yuri Petrovich or his friends.

Here is the list of 76 participants: K. Abe (Matsumoto, Japan), P. M. Akhmetjev (Moscow, Russia), S. V. Allyonov (Kolomna, Russia), V. A. Artamonov (Moscow, Russia), A. Bak (Bielefeld, Germany), V. V. Balashchenko (Minsk, Belarus), P. Baum (State College, USA), V. V. Belokurov (Moscow, Russia), B. Bojarski (Warsaw, Poland), N. Bokan (Belgrade, Serbia), D. Burghilea (Columbus, USA), O. M. Davydov (Chelyabinsk, Russia), M. Djorić (Belgrade, Serbia), Č. Dolićanin (Novi Pazar, Serbia), B. M. Doubrov (Minsk, Belarus), J. Eichhorn (Greifswald, Germany), A. Ermolitski (Minsk, Belarus), A. V. Ershov (Moscow, Russia), M. Frank (Leipzig, Germany), G. Garkusha (Trieste, Italy), Yu. E. Gliklikh (Voronezh, Russia), G. Guzhvina (Münster, Germany), B. Hajduk (Wroclaw, Poland), A. Ya. Helemsky (Moscow, Russia), A. A. Irmatov (Moscow, Russia), V. Kasimov (Baku, Azerbaijan), G. G. Kasparov (Nashville, USA), B. P. Komrakov (Leuven, Belgium), Yu. A. Kordyukov (Ufa, Russia), R. Krasauskas (Vilnius, Lithuania), A. Kuku (Berkeley, USA), S. V. Lapin (Saransk, Russia), R. Leandre (Dijon, France), V. P. Lexin (Kolomna, Russia), A. Lipkovski (Belgrade, Serbia), G. L. Litvinov (Moscow, Russia), M. A. Malakhaltsev (Kazan, Russia), V. M. Manuilov (Moscow, Russia), S. A. Melikhov (Moscow, Russia), D. V. Millionshchikov (Moscow, Russia), A. S. Mishchenko (Moscow, Russia), P. Mormul (Warsaw, Poland), E. A. Morozova (Moscow, Russia), Yu. V. Muranov (Vitebsk, Belarus), O. R. Musin (Moscow, Russia), V. M. Nezhinskij (St. Petersburg, Russia), I. M. Nikonov (Moscow, Russia), T. E. Panov (Moscow, Russia), D. Perrot (Lyon, France), S. S. Podkorytov (St. Petersburg, Russia), F. Yu. Popelensky (Moscow, Russia), P. S. Popov (Moscow, Russia), Z. Rakić (Belgrade, Serbia), A. Yu. Savin (Moscow, Russia), A. G. Sergeev (Moscow, Russia), V. V. Sharko (Kiev, Ukraine), G. I. Sharygin (Moscow, Russia), E. T. Shavgulidze (Moscow, Russia), A. I. Shtern (Moscow, Russia), M. I. Shtogrin (Moscow, Russia), A. B. Skopenkov (Moscow, Russia), M. B. Skopenkov (Moscow, Russia), A. B. Sossinski (Moscow, Russia), A. N. Starkov (Moscow, Russia), B. Yu. Sternin (Moscow, Russia), S. Terzić (Podgorica, Montenegro), A. Thom (Münster, Germany), E. V. Troitsky (Moscow, Russia), A. Yu. Volovikov (Moscow, Russia), S. Vukmirović (Belgrade, Serbia), A. V. Zarelua (Moscow, Russia), N. I. Zhukova (Nizhnij Novgorod, Russia), Yu. Y. Zhuraev (Samarkand, Uzbekistan), N. Yu. Zhuraeva (Samarkand, Uzbekistan), V. Zoller (Budapest, Hungary), V. G. Zvjagin (Voronezh, Russia).

The title of the conference demonstrates the expansiveness of Yu. P. Solovyov’s scientific interests — from classical topology to mathematical physics.

Initially, the research work of Yuri Petrovich was related to that of his adviser, Alexandr Sergeevich Mishchenko. Solovyov’s first result was obtained as early as in 1975, when he was a student, in the theory of elliptic operators, which was flourishing at that time. In topology, the necessity of studying the so-called Fredholm representations of discrete groups appeared. Properties of Fredholm operators realized as elliptic differential or pseudodifferential operators on compact manifolds were already well understood, but no nontrivial (from the point of view of discrete groups) examples of Fredholm operators were known, as an infinite discrete group cannot act freely on a compact manifold. Yu. P. Solovyov suggested a good, original idea: consider a differential elliptic operator in a Euclidean space, which is noncompact, and

Translated from *Fundamentalnaya i Prikladnaya Matematika*, Vol. 13, No. 7, pp. 3–6, 2007.

study its behavior with respect to the action of the free Abelian group of translations. As a result, he obtained a Fredholm representation of a free Abelian group with a nontrivial Chern character, providing, thus, a simple proof of the Novikov conjecture for free Abelian groups.

Then in 1975 he succeeded in generalizing this approach to more complicated spaces, the so-called Bruhat–Tits buildings. This result implies the proof of the Novikov conjecture for a wide class of discrete groups, namely, for lattices in linear algebraic groups over local, locally compact fields. This result was so strong that up till now it has only been slightly strengthened.

Not only these two papers, but a whole series was devoted to a systematic study of topological problems related to the Novikov conjecture on homotopy invariance of higher signatures. In particular, he discovered that higher signatures can be defined not only for smooth manifolds, but also for rationally homological ones. This possibility was based on the analysis of the construction of Hermitian K-theory using the so-called algebraic Poincaré complexes, which generalize the Poincaré duality of combinatorial manifolds from one side and Hermitian quadratic forms from the other side. The theory of sheaves of algebraic Poincaré complexes constructed jointly with A. S. Mishchenko led to a purely homotopic proof of the Novikov conjecture for those cohomology classes that can be obtained from Fredholm representations, and also to Hirzebruch-type formulas for arbitrary C^* -algebras.

Another problem of interest to Solovyov concerned the relation between the Novikov conjecture and the Baum–Connes conjecture. Here, the problem was essentially reduced to the description of elements of Hermitian K-theory that can be obtained from closed combinatorial manifolds. Yu. P. Solovyov gave this description in purely homotopical terms in 1976.

His further study of Hermitian K-theory in 1978–1982 led to interesting results on its classifying space and the corresponding homotopic multiplication, on the correlation between different approaches to the definition of Hermitian K-theory. In particular, he proved in 1982 that the analogs of the algebraic K-theory of Volodin–Wagoner constructed from the root systems B_n , C_n , and D_n are equivalent to Hermitian algebraic K-theory. He described the homotopic structure of the part of the Wall groups that comes from signatures. A very important result in this research was his construction of the Hermitian analog $KU(X)$ of the Waldhausen functor for topological spaces.

A series of papers by Yuri Petrovich with his students (R.-L. Krasauskas, S. V. Lapin, V. A. Kolosov) was devoted to homology theories with inner symmetries. It was shown in 1987 that such theories are defined by the so-called twisted simplicial groups, each of which can be uniquely presented as an extension of one of the seven simple twisted simplicial groups by an appropriate simplicial group. The deepest and most important results were obtained by Yu. P. Solovyov for dihedral homology, which is one of the seven basic types of homology with inner symmetries.

It turned out that dihedral homology is closely related to Hermitian K-theory. Informally speaking, dihedral homology is the logarithm of algebraic Hermitian K-theory in some sense. Yu. P. Solovyov discovered one more aspect of this relation that uses a Hermitian analog for the K-functor of Waldhausen, which is a generalization by Solovyov of Hermitian K-theory to topological spaces.

Yu. P. Solovyov applied dihedral homology to calculate invariants of the space of homotopic equivalences and the space of homeomorphisms of combinatorial manifolds. He showed that the calculation of rational homotopy groups for the group of homeomorphisms of a simply connected topological space reduces to the calculation of a dihedral homology of some special objects, which, in turn, can be evaluated by the minimal-model method of Sullivan. These results allowed one to calculate rational homotopy groups for the groups of homeomorphisms of complex and quaternion projective spaces and of some Grassmann manifolds.

In another large series of papers, he jointly with co-authors (V. V. Belokurov and E. T. Shavgulidze) developed a general approach to construction of absolutely converging series for approximation (with arbitrary precision) of functional integrals appearing in quantum physics, nonlinear differential equations theory, and topology. Traditionally, such functionals are evaluated by methods of perturbation theory, where series are asymptotic power decompositions of integrals with respect to some parameter. Despite the good accordance of the sum of the series' first several elements with experimental data for physical

problems with small parameter values, in general the question on the possibility of approximation for the corresponding functional integrals (with arbitrary precision) by a finite number of a divergent series is open. Within the framework of the approach suggested by Yu. P. Solov'yov and co-authors, it turned out to be possible to evaluate functional integrals with any precision by summing a finite number of elements of an absolutely converging series of a new perturbation theory. In particular, indices that describe the behavior of the thermal capacity of helium near the critical point were calculated with unprecedented precision. This method was also applied to generalized summation of a wide class of divergent series that can be represented as Taylor series for smooth completely monotonic functions. In this case, the sum of the series is defined as the value of the function at a given point. Using this approach, one can evaluate with any precision the sum of some divergent series, which cannot be calculated by other methods.

The conference talks covered all directions of research led by Yu. P. Solov'yov during his life. Not all of the talks are presented here. Some of them will be published in the *Journal of K-Theory*.

The papers can be divided into several directions of research.

- (1) Classical problems of topology (P. Akhmetiev, S. Lapin, I. Nikonov, A. Ershov).
- (2) Symmetries (V. Balashchenko, A. Volovikov).
- (3) Equations on manifolds (Yu. Gliklikh, N. Zhuraeva).
- (4) Functional analysis and C^* -algebras (A. Helemskii, R. Leandre, A. Pavlov, E. Troitsky, T. Shulman).
- (5) Group representations and ergodic theory (A. Troitskaya, A. Shtern).

V. M. Manuilov

Faculty of Mechanics and Mathematics, Moscow State University, Moscow 119991 GSP, Russia

A. S. Mishchenko

Faculty of Mechanics and Mathematics, Moscow State University, Moscow 119991 GSP, Russia