ERRATUM

Erratum to: Application of isothermal titration calorimetry for evaluation of water–acetone and water–dimethylsulfoxide solvents influence on the molecular complex formation between 18-crown-6 and triglycine at 298.15 K

T. R. Usacheva¹ · L. Pham Thi¹ · I. V. Terekhova² · R. S. Kumeev² · V. A. Sharnin^{1,2}

Received: 20 May 2016 / Accepted: 21 May 2016 / Published online: 13 June 2016
© Akadémiai Kiadó, Budapest, Hungary 2016

Erratum to: J Therm Anal Calorim (2015)
121:975–981
DOI 10.1007/s10973-015-4630-0

The correct numeration for H-1 and H-3 peaks is as follows (Fig. 2).

Fig. 2 ^1H NMR spectra of 3Gly ($0.001 \text{ mol kg}^{-1}$) in water and solution of 18C6 ($0.165 \text{ mol kg}^{-1}$) at $T = 298.15 \text{ K}$

The online version of the original article can be found under doi:[10.1007/s10973-015-4630-0](https://doi.org/10.1007/s10973-015-4630-0).

✉ T. R. Usacheva
oxt@isuct.ru

¹ Research Institute of Thermodynamics and Kinetics of Chemical Processes, Ivanovo State University of Chemistry and Technology, pr. Sheremetevski 7, 153000 Ivanovo, Russia

² G.A. Krestov Institute of Solution Chemistry of the Russian Academy of Sciences, Akademicheskaya str. 1, 153045 Ivanovo, Russia