Strategies to Reduce Alcohol-Exposed Pregnancies

R. Louise Floyd · Shahul Ebrahim · James Tsai ·
Mary O'Connor · Robert Sokol

Published online: 8 March 2007
© Springer Science+Business Media, LLC 2007

Erratum to: Maternal and Child Health Journal, Vol. 10,
Supplement 7, September 2006 **DOI: 10.1007/s10995-006-
0116-9.**

This is how the correct footnote should read:
Frequent refers to ≥ 7 drinks/week or binge. Binge drink-
ing refers to ≥ 5 drinks on one occasion.

The online version of the original article can be found at:
<http://dx.doi.org/10.1007/s10995-006-0116-9>

R. L. Floyd · S. Ebrahim · J. Tsai
Centers for Disease Control & Prevention,
Atlanta, Georgia

M. O'Connor
Department of Psychiatry and Biobehavioral Sciences, University
of California at Los Angeles, David Geffen School of Medicine,
Los Angeles, California

R. Sokol
Department of Obstetrics and Gynecology, Wayne State
University, School of Medicine,
Detroit, Michigan

S. Ebrahim (✉)
Centers for Disease Control & Prevention (E86),
Atlanta, Georgia 30333
e-mail: sebrahim@cdc.gov