CK2 mediates phosphorylation and ubiquitin-mediated degradation of the PML tumor suppressor

P. P. Scaglioni · T. M. Yung · S. Choi · C. Baldini ·
G. Konstantinidou · P. P. Pandolfi

Published online: 6 May 2009
© Springer Science+Business Media, LLC. 2009

Erratum to: Mol Cell Biochem (2008) 316:149–154
DOI: 10.1007/s11010-008-9812-7

In the paper, the name of the third author is given as S. C. Choi. Dr. Sooyeon Choi does not have a middle initial. Her correct name should read “S. Choi”.

Everything else in the paper remains correct.

The online version of the original article can be found under doi:
[10.1007/s11010-008-9812-7](https://doi.org/10.1007/s11010-008-9812-7).

P. P. Scaglioni (✉) · S. Choi · C. Baldini · G. Konstantinidou
Division of Hematology-Oncology, University of Texas
Southwestern Medical Center, Dallas, TX 75390-8852, USA
e-mail: pier.scaglioni@utsouthwestern.edu

T. M. Yung
Cancer Biology and Genetics Program, Sloan-Kettering Institute,
Memorial Sloan-Kettering Cancer Center, New York,
NY 10021, USA

P. P. Pandolfi
Cancer Genetics Program, Beth Israel Deaconess Cancer Center,
Harvard Medical School, Boston, MA 02215, USA

P. P. Pandolfi
Department of Medicine, Beth Israel Deaconess Medical Center,
Harvard Medical School, Boston, MA 02215, USA