The dose–volume interaction in adult supratentorial low-grade glioma: higher radiation dose is beneficial among patients with partial resection

C. Leighton · B. Fisher · D. Macdonald · L. Stitt ·
G. Bauman · J. Cairncross

Published online: 10 July 2007
© Springer Science+Business Media B.V. 2007

Erratum to: J Neurooncol (2007) 82:165–170
DOI 10.1007/s11060-006-9141-1

The original version of this article unfortunately contained a mistake. The sequence of the author names was incorrect.

The correct sequence is

C. Leighton, B. Fisher, D. Macdonald, L. Stitt,
G. Bauman, J. Cairncross

The online version of the original article can be found under doi:
[10.1007/s11060-006-9141-1](https://doi.org/10.1007/s11060-006-9141-1).

C. Leighton · B. Fisher (✉) · G. Bauman
Division of Radiation Oncology, London Regional Cancer
Centre Program, London Health Sciences Hospital, London,
Ontario, Canada
e-mail: barb.fisher@lrcc.on.ca

D. Macdonald
Department of Medical Oncology, London Regional Cancer
Program and Department of Neurooncology, University
of Western Ontario, London, Ontario, Canada

L. Stitt
Department of Epidemiology and Biostatistics, University
of Western Ontario and Clinical Research Unit, London
Regional Cancer Program, London Health Sciences Hospital,
London, Ontario, Canada

J. Cairncross
Tom Baker Cancer Centre, Calgary, Alberta, Canada