Erratum to: Correlating the Metabolic Stability of Psychedelic 5-HT_{2A} Agonists with Anecdotal Reports of Human Oral Bioavailability

Sebastian Leth-Petersen · Christoffer Bundgaard ·
Martin Hansen · Martin A. Carnerup ·
Jan Kehler · Jesper Langgaard Kristensen

Published online: 28 March 2014
© Springer Science+Business Media New York 2014

Erratum to: Neurochem Res DOI 10.1007/s11064-014-1253-y

The presentation of Table 1 was published incorrectly due to an inadvertent error during typesetting. The corrected Table 1 is given below. We apologize for the technical error.

Table 1 Hepatic stability of psychedelic 2C-X's (**1a–10a**) and *N*-benzyls (**1b–10b** and **1c–10c**) given as intrinsic clearance (CL_{int} in L/kg/h)

R =	CL _{int} (L/Kg/h)			
I	1a: 0.20	1b: 4.1	1c: 9.9	
CH ₃	2a: 0.27	2b: 4.4	2c: 6.3	
SEt	3a: 0.48	3b: 19	3c: 42	
Br	4a: 0.51	4b: 3.3	4c: 5.4	
CN	5a: 0.58	5b: 5.0	5c: 7.6	
SPr	6a: 0.61	6b: 13	6c: 36	
CF ₃	7a: 0.62	7b: 8.8	7c: 8.4	
SMe	8a: 0.65	8b: 12	8c: 29	
Cl	9a: 0.74	9b: 2.9	9c: 5.8	
H	10a: 8.9	10b: 14	10c: 16	
Average	0.5	9	17	
	(exluding 10a)			

The compounds are listed according to the metabolic rate of the 2C-X series

The online version of the original article can be found under doi:10.1007/s11064-014-1253-y.

S. Leth-Petersen · M. Hansen · J. L. Kristensen (✉)
Department of Drug Design and Pharmacology, Faculty of
Health and Medical Sciences, University of Copenhagen,
Universitetsparken 2, 2100 Copenhagen, Denmark
e-mail: jesper.kristensen@sund.ku.dk

C. Bundgaard · M. A. Carnerup · J. Kehler
Department of Discovery Chemistry and DMPK, H. Lundbeck
A/S, Ottiliavej 9, 2500 Valby, Denmark