ERRATUM

Erratum to: Exogenous Adipokine Peptide Resistin Protects Against Focal Cerebral Ischemia/Reperfusion Injury in Mice

Jiangtao Zhu¹ · Di Wu¹ · Chenyu Zhao¹ · Man Luo¹ · Ronald C. Hamdy³ · Balvin H. L. Chua³ · Xingshun Xu^{1,2} · Zhigang Miao²

Published online: 25 July 2017
© Springer Science+Business Media, LLC 2017

Erratum to: Neurochem Res
DOI 10.1007/s11064-017-2326-5

Fig. 5A (ATP level) is wrong. The corrected Fig. 5 is given below.

The original version of this article unfortunately contained a mistake. The presentation of x-axis coordinates of

Fig. 5 Resistin treatment increased cellular ATP level after MCAO. ATP levels in ischemic brain tissues were measured at 24 h after MCAO (a). The food intake was recorded at 1, 3, and 7 days after

MCAO (b). Serum glucose was also measured at 1, 3, and 7 days after MCAO (c). N = 6, *P < 0.05, versus sham group; #P < 0.05, versus I/R + R group

The online version of the original article can be found under doi:10.1007/s11064-017-2326-5.

✉ Zhigang Miao
sxmiaomiao@163.com

Xingshun Xu
Xinshunxu@suda.edu.cn

¹ Department of Neurology, The Second Affiliated Hospital of Soochow University, Suzhou City, China

² Institute of Neuroscience, Soochow University, Suzhou City, Jiangsu, China

³ Cecile Cox Quillen Laboratory of Geriatrics, James H. Quillen College of Medicine, East Tennessee State University, Johnson City, TN 37614, USA