Correction to: Chronic Administration of S-Allylcysteine Activates Nrf2 Factor and Enhances the Activity of Antioxidant Enzymes in the Striatum, Frontal Cortex and Hippocampus

Úrzula Franco-Enzástiga¹ · Ricardo A. Santana-Martínez¹ · Carlos A. Silva-Islas¹ · Diana Barrera-Oviedo² · María Elena Chánez-Cárdenas¹ · Perla D. Maldonado¹

Published online: 26 December 2017
© Springer Science+Business Media, LLC, part of Springer Nature 2017

Correction to:
Neurochemical Research (2017) 42:3041–3051
<https://doi.org/10.1007/s11064-017-2337-2>

The original version of this article unfortunately contained a mistake. In Fig. 5, the image corresponding to hippocampus with SAC 200 mg/kg is incorrect. The corrected Fig. 5 is given below. The figure caption corresponding to Fig. 5 is the same.

The original article can be found online at <https://doi.org/10.1007/s11064-017-2337-2>.

✉ Perla D. Maldonado
maldonado.perla@gmail.com

¹ Laboratorio de Patología Vascular Cerebral, Instituto Nacional de Neurología y Neurocirugía, Insurgentes Sur 3877, 14269 Mexico City, Mexico

² Departamento de Farmacología, Facultad de Medicina, Universidad Nacional Autónoma de México, 04510 Mexico City, Mexico

Fig. 5 Effect of chronic administration of SAC on 8-hydroxy-2-deoxyguanosine (8-OHdG) levels in the striatum, frontal cortex, and hippocampus (CA3). SAC (25, 50, 100 and 200 mg/kg, i.g.) was administered each 24 h during 90 days. Control (CT) group received saline solution in the same conditions. **a** Representative microphoto-

graphs of 8-OHdG immunodetection on each group are shown. White arrows immunoreactive cell. Magnification $\times 40$. **b** Graph represents the quantitative analysis scored in the different groups. Data are expressed as mean $\pm$ SEM of three animals per group