ERRATUM

Erratum to: Do chronic aboveground O₃ exposure and belowground pathogen stress affect growth and belowground biomass partitioning of juvenile beech trees (*Fagus sylvatica* L.)?

J. Barbro Winkler · Frank Fleischmann ·
Sebastian Gayler · Hagen Scherb ·
Rainer Matyssek · Thorsten E. E. Grams

Published online: 12 September 2009
© Springer Science + Business Media B.V. 2009

Erratum to: Plant Soil (2009) 323:31–44 DOI 10.1007/s11104-009-9968-2

Owing to an error during the submission process, the original article was inadvertently published with an incorrect version of Fig. 4. The correct image and caption is shown here.

Responsible Editor: Peter A.H. Bakker.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s11104-009-9968-2>.

J. B. Winkler (✉)
Institute of Soil Ecology, Department of Environmental
Engineering, Helmholtz Zentrum München,
German Research Center for Environmental Health,
Ingolstädter Landstr. 1,
85764 Neuherberg, Germany
e-mail: bwinkler@helmholtz-muenchen.de

F. Fleischmann
Pathology of Woody Plants, Department of Ecology
& Ecosystem Sciences, Center of Life and Food Science
Weihenstephan, Technische Universität München,
Am Hochanger 13,
85354 Freising, Germany

S. Gayler
Institute of Soil Ecology, Helmholtz Zentrum München,
German Research Center for Environmental Health,
Ingolstädter Landstr. 1,
85764 Neuherberg, Germany

H. Scherb
Institute of Biomathematics and Biometry,
Helmholtz Zentrum München,
German Research Center for Environmental Health,
Ingolstädter Landstr. 1,
85764 Neuherberg, Germany

R. Matyssek · T. E. E. Grams
Ecophysiology of Plants,
Department of Ecology & Ecosystem Sciences,
Center of Life and Food Science Weihenstephan,
Technische Universität München,
Am Hochanger 13,
85354 Freising, Germany

Fig. 4 Total tree biomass of tree growing under ambient O₃ (white bars) or elevated O₃ (grey bars) and on non-inoculated lysimeters (plain bars) or on lysimeters with *P. citricola* infection (striped bars). The upper part of the figure represents the aboveground biomass and the lower part the root mass. Given are means $\pm$ standard error ($n=8$)