Erratum to: Analysis of the *Wsi18*, a stress-inducible promoter that is active in the whole grain of transgenic rice

Nari Yi · Se-Jun Oh · Youn Shic Kim ·
Hyun-Jun Jang · Su-Hyun Park · Jin Seo Jeong ·
Sang Ik Song · Yang Do Choi · Ju-Kon Kim

Published online: 31 August 2010
© Springer Science+Business Media B.V. 2010

Erratum to: Transgenic Res DOI 10.1007/s11248-010-9400-y

Due to an unfortunate mistake, the incorrect figure caption of Fig. 3 has been used in the above-

mentioned publication. The correct representation of Fig. 3 and its caption are published on the following page and should be treated as definitive by the reader.

The online version of the original article can be found under doi:[10.1007/s11248-010-9400-y](https://doi.org/10.1007/s11248-010-9400-y).

N. Yi · S.-J. Oh · Y. S. Kim · H.-J. Jang · S.-H. Park ·
J. S. Jeong · S. I. Song · J.-K. Kim (✉)
School of Biotechnology and Environmental Engineering,
Myongji University, Yongin 449-728, Korea
e-mail: jukon306@gmail.com

N. Yi · Y. D. Choi
School of Agricultural Biotechnology, Seoul National
University, Seoul 151-921, Korea

S.-J. Oh
Syngenta Seeds Co., Ltd, 100 Gongpyeong-dong,
Seoul 110-702, Korea

Fig. 3 Activity of the *Wsi18* promoter in different tissues and growth stages of transgenic rice plants (line 2 shown in Fig. 2b). **a** GFP fluorescence in transgenic calli. GFP fluorescence was detected in calli exposed to drought, NaCl (400 mM), and ABA (100 μ M) for the indicated time points using a luminescent-image analyzer. NT non-transgenic calli, *OsCc1* calli induced from the *OsCc1*::GFP transgenic seeds (Jang et al. 2002), *Wsi18* calli induced from the *Wsi18*::GFP transgenic seeds. Bar = 1 mm. **b** Confocal microscopic images of GFP fluorescence in leaf and root tissues of transgenic plants. GFP fluorescence was detected in a leaf, a root apex, and a elongating region of a root of the *Wsi18*::GFP and *OsCc1*::GFP plants before (normal) and after exposure to drought, NaCl (400 mM), and ABA (100 μ M) for 2–6 h.

The images collected in the *green* and *i* channels of the confocal microscope were merged. Bars = 1 mm. S stomata. **c** GFP fluorescence in whole bodies of the *Wsi18*::GFP and *OsCc1*::GFP plants. GFP fluorescence was detected in 6-day-old etiolated plants exposed to drought, NaCl (400 mM), and ABA (100 μ M) for the indicated time points using a luminescent-image analyzer. Bar = 1 cm. **d** GFP fluorescence in transgenic flowers. GFP fluorescence was detected, by use of a stereomicroscope, in a whole spikelet (left panel) and floral organs inside a spikelet (right panel) at the stage of meiosis before (Normal) and after drought treatment. For drought treatment, water was withheld from plants in pots, in a greenhouse, for three days. An, anther; F, filament; Lm, lemma; Lo, lodicule; P, palea. Bars = 1 mm