Erratum to: Characterization of the spectrum of insecticidal activity of a double-stranded RNA with targeted activity against Western Corn Rootworm (*Diabrotica virgifera virgifera* LeConte)

Pamela M. Bachman · Renata Bolognesi · William J. Moar · Geoffrey M. Mueller · Mark S. Paradise · Parthasarathy Ramaseshadri · Jianguo Tan · Joshua P. Uffman · JoAnne Warren · B. Elizabeth Wiggins · Steven L. Levine

Published online: 18 September 2013
© Springer Science+Business Media Dordrecht 2013

Erratum to: Transgenic Res DOI 10.1007/s11248-013-9716-5

Due to an unfortunate turn of events, an error was introduced in Table 1 of the above-mentioned publication. Two LC₅₀ values in the first two adjacent rows were inadvertently transposed. Furthermore, in the

paragraph just above Figure 1, the words ‘control’ and ‘test’ were interchanged. Therefore, the sentence must read: “Additionally, there was no significant effect on CPB growth with mean test and control weights of 8.5 ± 0.2 and 8.2 ± 0.4 mg, respectively ($p > 0.05$).”

The correct table is published on the next page and should be treated as definitive by the reader.

The online version of the original article can be found under doi:[10.1007/s11248-013-9716-5](https://doi.org/10.1007/s11248-013-9716-5).

P. M. Bachman (✉) · W. J. Moar · G. M. Mueller · M. S. Paradise · J. Tan · J. P. Uffman · J. Warren · S. L. Levine
Monsanto Company, 800 N Lindbergh Blvd., St. Louis, MO 63167, USA
e-mail: pamela.m.bachman@monsanto.com

R. Bolognesi · P. Ramaseshadri · B. Elizabeth Wiggins
Monsanto Company, 700 Chesterfield Parkway W, Chesterfield, MO 63017, USA

Table 1 Bioassay results from direct feeding studies with DvSnf7 dsRNA using insects from four Orders representing ten insect families

Order	Family	Subfamily	Species	Bioassay duration (days)	End points	LC ₅₀ ^a or no observed effect concentration ^b (ng/mL or g diet)
Coleoptera	Chrysomelidae	Galerucinae	<i>D. virgifera virgifera</i>	12	Survival	4.4 ^a
	Chrysomelidae	Galerucinae	<i>D. undecimpunctata howardi</i>	12	Survival	1.2 ^a
	Chrysomelidae	Chrysomelinae	<i>L. decemlineata</i>	12	Survival, Growth	5,000 ^b
	Tenebrionidae	Tenebrioninae	<i>T. castaneum</i>	30	Survival, Growth	5,000 ^b
	Coccinellidae	Coccinellinae	<i>C. maculata</i>	24	Survival, Growth, Development	3,000 ^b
	Coccinellidae	Epilachninae	<i>E. varivestis</i>	28	Survival, Growth, Development	3,000 ^b
	Carabidae	Harpalinae	<i>P. chalcites</i>	35	Survival, Growth, Development	5,000 ^b
Hemiptera	Anthoridae	Anthocorinae	<i>O. insidiosus</i>	9	Survival, Growth, Development	5,000 ^b
Hymenoptera	Eulophidae	Entedoninae	<i>P. foveolatus</i>	21	Survival	3,000 ^b
	Pteromalidae	Pteromalinae	<i>N. vitripennis</i>	20	Survival	5,000 ^b
Lepidoptera	Noctuidae	Noctuinae	<i>S. frugiperda</i>	8	Survival, Growth	500 ^b
		Heliothinae	<i>H. zea</i>	12	Survival, Growth	5,000 ^b
	Crambidae	Pyraustinae	<i>O. nubilalis</i>	12	Survival, Growth	5,000 ^b
	Bombycidae	Bombycinae	<i>B. mori</i>	14	Survival, Growth	5,000 ^b

^a LC₅₀ data reported in Bolognesi et al. (2012)

^b No observed effect concentration is equal to the maximum concentration tested in these bioassays. Concentrations ranged from 500 to 5000 ng/mL or ng/g diet and represent 250–2,500 times the maximum expected environmental concentration of DvSnf7 based on greenhouse expression levels