A rare complication after circumcision: Keloid of the penis

Fikret Erdemir · Ozgur Gokce · Oner Sanli · Ates Kadioglu · Bekir Suha Parlaktas · Nihat Uluocak · Isin Kilicaslan

Published online: 6 July 2007
© Springer Science+Business Media B.V. 2007

Erratum to: Int Urol Nephrol (2006) 38(3–4):609–611
DOI 10.1007/s11255-006-0021-6

The information about the authors and their affiliations was incomplete. The correct list of authors and their affiliations are shown above and below.

Fikret Erdemir: Department of Urology, Istanbul Faculty of Medicine, Istanbul University, Istanbul, Turkey. E-mail: fikreterdemir@mynet.com

Ozgur Gokce: Department of Urology, Istanbul Faculty of Medicine, Istanbul University, Istanbul, Turkey

Oner Sanli: Department of Urology, Istanbul Faculty of Medicine, Istanbul University, Istanbul, Turkey

Ates Kadioglu: Department of Urology, Istanbul Faculty of Medicine, Istanbul University, Istanbul, Turkey

Bekir Suha Parlaktas: Department of Urology, Istanbul Faculty of Medicine, Istanbul University, Istanbul, Turkey

Nihat Uluocak: Department of Urology, Istanbul Faculty of Medicine, Istanbul University, Istanbul, Turkey

Isin Kilicaslan: Department of Pathology, Istanbul Faculty of Medicine, Istanbul University, Istanbul, Turkey

The online version of the original article can be found under doi: [10.1007/s11255-006-0021-6](https://doi.org/10.1007/s11255-006-0021-6).

F. Erdemir (✉) · O. Gokce · O. Sanli · A. Kadioglu · B. S. Parlaktas · N. Uluocak
Department of Urology, Istanbul Faculty of Medicine,
Istanbul University, Istanbul, Turkey
e-mail: fikreterdemir@mynet.com

I. Kilicaslan
Department of Pathology, Istanbul Faculty of Medicine,
Istanbul University, Istanbul, Turkey