Oxidative stress markers and C-reactive protein in end-stage renal failure patients on dialysis

**Elisabeth C. Samouilidou · Eirini J. Grapsa ·
Ioannis Kakavas · Antonios Lagouranis ·
Basilis Agrogiannis**

Published online: 27 September 2007
© Springer Science+Business Media B.V. 2007

**Erratum to: Int Urol Nephrol (2003) 35:393–397
DOI 10.1023/B:UROL.0000022846.83505.3f**

In the article “Oxidative stress markers and C-reactive protein in end-stage renal failure patients on dialysis” by Elisabeth C. Samouilidou, Eirini J. Grapsa, Ioannis Kakavas, Antonios Lagouranis and Basilis Agrogiannis, published in International

Urology and Nephrology-issue 35(3), pages 393–397, Fig. 2 was incorrect. Please find the correct figure below.

The online version of the original article can be found under doi: [10.1023/B:UROL.0000022846.83505.3f](https://doi.org/10.1023/B:UROL.0000022846.83505.3f)

E. C. Samouilidou
Biochemical Department, Athens, Greece

E. J. Grapsa (✉) · A. Lagouranis
Renal Unit ‘Alexandra’ Hospital, 80, Vas. Sophias Str.,
Athens 11528, Greece
e-mail: egrapsa@teledomenet.gr

I. Kakavas
Renal Unit ‘Ippokration’ Hospital, Athens, Greece

B. Agrogiannis
Renal Unit ‘Areteion’ Hospital, Athens, Greece

Fig. 2 Plasma CRP levels in healthy individuals (NC), peritoneal dialysis (PD) and hemodialysis (HD) patients before and after dialysis. Columns represent mean values $\pm$ S.E.M. Significance: $p < 0.001$ in HD patients before or after dialysis vs. NC and PD, $p < 0.05$ in PD patients vs. NC