Effect of tephra depth on vegetation development in areas affected by volcanism

Mariela Gómez-Romero · Roberto Lindig-Cisneros ·
Sebastiana Galindo-Vallejo

Published online: 28 February 2007
© Springer Science + Business Media B.V. 2007

Erratum to: Plant Ecol DOI 10.1007/s11258-005-9017-z

This erratum is published as wrong table & figure legends were published earlier. The re-corrected legends should be now read as:

Table 2 Total area, number of vegetation patches, and canopy cover of all vegetation patches in relation with tephra depth class in the “Mesa de Cutzato” 1.44 ha study plot

Fig. 3 Relationships between tephra depth (as mean depth for each depth class) and the density of vegetation patches in “a” (calculated as the

number of patches per hectare) ($R^2 = 0.67$, $F_{(1,6)} = 12.08$, $P = 0.013$) and with vegetation patch area per hectare in “b”

Fig. 5 Relationships between depth of the tephra layer and plant height and depth of the tephra layer and biomass (as dry weight), for *Lupinus elegans* plants growing on fine grained tephra (a) and coarse grained tephra (b) under greenhouse conditions. Two plants growing directly on the growing medium (treatment 1) were damaged by an un-identified herbivore and therefore excluded from the regression analysis (indicated by triangles in graph a)

The online version of the original article can be found at
<http://dx.doi.org/10.1007/s11258-005-9017-z>

M. Gómez-Romero · R. Lindig-Cisneros ·
S. Galindo-Vallejo
Laboratorio de Ecología de Restauración, Facultad
de Biología, Universidad Michoacana de San Nicolás
de Hidalgo, Ciudad Universitaria, Morelia,
Michoacán, México

R. Lindig-Cisneros (✉)
Apartado Postal 18, Administración 3, Santa Maria,
Morelia C.P. 58091 Michoacán, México
e-mail: lindig@zeus.umich.mx