Erratum

A Parasitological Survey of European Sea Bass (*Dicentrarchus Labrax*) and Gilthead Sea Bream (*Sparus Aurata*) Cultured in Italy

M.L. Fioravanti*, M. Caffara, D. Florio, A. Gustinelli and F. Marcer

Department of Veterinary Public Health and Animal Pathology, Faculty of Veterinary Medicine, University of Bologna, Ozzano Emilia (BO), Italy

*Correspondence: Dipartimento di Sanità Pubblica Veterinaria e Patologia Animale – Facoltà di Medicina Veterinaria, via Tolara di Sopra, 50 – 40064, Ozzano Emilia (BO), Italy
E-mail: fioravanti@vet.unibo.it

Veterinary Research Communications, **30** (Suppl.1) 2006, 249–252

Tables I and II in this article were badly laid out, for which the publishers apologise. The correct layout of the Tables is given below:

TABLE I
Parasites found in European sea bass (*Dicentrarchus labrax*)

Parasites from European sea bass	No. of positive subjects – <i>percentage of positivity</i>					Total
	Extensive	Intensive			SC	
		I	IFC	OFC		
Sarcomastigophora						
<i>Amyloodinium ocellatum</i>	99–34.1%	52–12.6%	–	–	–	151–11.6%
<i>Cryptobia branchialis</i>	45–15.5%	2–0.5%	–	–	1–0.5%	48–3.7%
Ciliophora						
<i>Trichodina</i> spp.	63–21.7%	55–13.3%		–	–	118–9.1%
Sessiline Peritrichs	3–1%	–	–	–	–	3–0.2%
Apicomplexa						
<i>Eimeria bouixi</i>	1–0.3%	8–1.9%	8–2.2%	6–15%	3–1.6%	26–2%
Microspora	–	28–6.8%	–	–	–	28–2.2%
Myxozoa						
<i>Sphaerospora dicentrarchi</i>	139–47.9%	243–59%	142–38.7%	13–32.5%	5–2.6%	542–41.6%
<i>S. testicularis</i>	–	–	17–4.6%	–	–	17–1.3%
<i>Ceratomyxa labracis</i>	2–0.7%	34–8.3%	7–1.9%	6–15%	18–9.3%	67–5.1%
<i>C. diplodae</i>	–	7–1.7%	30–8.2%	–	–	37–2.8%
<i>Enteromyxum leei</i>	–	15–3.6%	–	–	–	15–1.2%
Platyhelminthes Monogenea						
<i>Diplectanum aequans</i>	138–47.6%	221–53.6%	142–38.7%	1–2.5%	13–6.7%	515–39.6%
Arthropoda Crustacea						
<i>Caligus minimus</i>	–	4–1%	–	–	–	4–0.3%

The online version of the original article can be found at <http://dx.doi.org/10.1007/s11259-006-0053-5>

TABLE II
Parasites found in gilthead sea bream (*Sparus aurata*)

Parasites from gilthead sea bream	No. positive subjects – percentage of positivity					Total
	Extensive	Intensive				
		I	IFC	OFC	SC	
Sarcomastigophora						
<i>Amyloodinium ocellatum</i>	–	10–5.8%	–	–	–	10–2.2%
Scuticociliatida	–	5–2.9%	3–2.8%	–	–	8–1.7%
Ciliophora						
<i>Cryptocaryon irritans</i>	–	22–12.7%	–	–	–	22–4.8%
<i>Trichodina</i> spp.	10–30.3%	–	2–1.9%	–	–	12–2.6%
Sessiline Peritrichs	16–48.5%	–	11–10.4%	–	–	27–5.9%
Microspora	–	–	15–14.2%	–	–	15–3.3%
Myxozoa						
<i>Ceratomyxa sparusaurati</i>	–	3–1.7%	2–1.9%	–	–	5–1.1%
<i>Polysporoplasma sparis</i>	–	5–2.9%	48–45.3%	–	–	53–11.6%
<i>Enteromyxum leei</i>	–	20–11.6%	–	43–71.7%	–	63–13.8%
<i>Leptotheca sparidarum</i>	–	32–18.5%	–	–	–	32–7%
Platyhelminthes Monogenea						
<i>Sparicotyle chrysophrii</i>	–	–	13–12.3%	–	15–17.4%	28–6.1%
<i>Furnestinia echeneis</i>	–	5–2.9%	7–6.6%	–	–	12–2.6%
Platyhelminthes Digenea						
Sanguinicolidae (eggs)	–	–	12–11.3%	–	–	12–2.6%
Arthropoda Crustacea						
<i>Caligus</i> spp.	3–9.1%	–	–	–	–	3–0.7%
<i>Ergasilus</i> spp.	–	–	12–11.3%	–	–	12–2.6%