Antibacterial activity and phytochemical evidence for the plant origin of Turkish propolis from different regions

Sibel Silici · Mehmet Ünlü · Gülhan Vardar-Ünlü

Published online: 11 November 2008
© Springer Science+Business Media B.V. 2008

Erratum to: World J Microbiol Biotechnol (2007) 23:1797–1803
DOI 10.1007/s11274-007-9430-7

Due to unfortunate circumstances, errors were introduced in the above-mentioned article. The correct text is published below and should be treated as definitive by the reader:

Abstract, page 1797. The fourth sentence should read: “One of the main components of *Eucalyptus* propolis were aromatic acids, mainly cinnamic acid and its esters, that are usually found in *Eucalyptus* species resins.”

Page 1800, the first sentence of the last paragraph should read: “Our investigation showed that one of the main

components of *Eucalyptus* propolis were aromatic acids, such as cinnamic acid and its esters (Table 1) that are usually found in *Eucalyptus* species resins.”

Table 1, page 1800. The first two italicized headings must be deleted and replaced by a single heading relating to items 1–17 and should read:

Phosphoric and carboxylic acids

The online version of the original article can be found under doi:[10.1007/s11274-007-9430-7](https://doi.org/10.1007/s11274-007-9430-7).

S. Silici (✉)
Department of Animal Science, S. Çıkıkcıoğlu Vocational
College, Erciyes University, 38039 Kayseri, Turkey
e-mail: silicis@erciyes.edu.tr

M. Ünlü · G. Vardar-Ünlü
Department of Microbiology, Faculty of Medicine, Cumhuriyet
University, Sivas 58140, Turkey