Antiviral activity of *Cleome rosea* extracts from field-grown plants and tissue culture-derived materials against acyclovir-resistant *Herpes simplex* viruses type 1 (ACVr-HSV-1) and type 2 (ACVr-HSV-2)

Claudia Simões · Tatiana Carvalho de Castro ·
Lívia da Silva Cordeiro · Norma Albarello ·
Elisabeth Mansur · Maria Teresa Villela Romanos

Published online: 18 September 2009
© Springer Science+Business Media B.V. 2009

Erratum to: World J Microbiol Biotechnol
DOI 10.1007/s11274-009-0147-7

Unfortunately, in the article “Antiviral activity of *Cleome rosea* extracts from field-grown plants and tissue culture-derived materials against acyclovir-resistant *Herpes simplex* viruses type 1 (ACVr-HSV-1) and type 2 (ACVr-HSV-2)”,

World J Microbiol Biotechnol, Table 3 has been incorrectly published.

The correct Table 3 is presented below:

The online version of the original article can be found under
doi:[10.1007/s11274-009-0147-7](https://doi.org/10.1007/s11274-009-0147-7).

C. Simões · T. C. de Castro · L. da Silva Cordeiro ·
N. Albarello · E. Mansur
Núcleo de Biotecnologia Vegetal, Instituto de Biologia Roberto
Alcantara Gomes, Universidade do Estado do Rio de Janeiro
(UERJ), Rio de Janeiro, RJ, Brazil

M. T. V. Romanos
Laboratório Experimental de Drogas Antivirais e Citotóxicas
(LEDAC), Departamento de Virologia, Instituto de
Microbiologia Prof. Paulo Góes, Universidade Federal do Rio de
Janeiro (UFRJ), Rio de Janeiro, RJ, Brazil

C. Simões (✉)
Departamento de Biologia Vegetal, Laboratório de
Biotecnologia de Plantas (LABPLAN), Instituto de Biologia
Roberto Alcantara Gomes, Universidade do Estado do Rio de
Janeiro, Centro Biomédico, Rua São Francisco Xavier, 524,
PHLC, Sala 509, Maracanã, CEP 20550-013 Rio de Janeiro, RJ,
Brazil
e-mail: labplan_uerj@yahoo.com.br

Table 3 Effective dose to reduce virus titers by 50% (EC_{50}) and selectivity index (SI) of *Cleome rosea* extracts

Extracts	CC_{50} ($\mu\text{g ml}^{-1}$)	ACVr-HSV-1		ACVr-HSV-2	
		EC_{50} ($\mu\text{g ml}^{-1}$)	SI	EC_{50} ($\mu\text{g ml}^{-1}$)	SI
In vivo plants (leaves) (MeOH)	68.7	12.9	5.3	9.3	7.4
Pigmented calluses (MeOH)	4972.2	54.2	91.7	52.4	94.8
Pigmented calluses (MeOH–HCl)	1535.2	43.8	35	62.6	24.5
Acyclovir	>200	>200	1	4.8	41.6

ACVr-HSV-1 acyclovir-resistant herpes simplex virus type 1, ACVr-HSV-2 acyclovir-resistant herpes simplex virus type 2, CC_{50} 50% cytotoxic concentration, EC_{50} effective concentrations to reduce virus titers by 50%, SI selectivity index, Acyclovir standard compound