Koichi Takahashi · Masaru Yogo · Satoshi Ishibashi

Stand development and regeneration during a 33-year period in a seral *Picea glehnii* forest, northern Japan

Published online: 16 December 2005
© The Ecological Society of Japan 2005

Ecol Res: DOI 10.1007/s11284-005-0093-2

When this article was published online, Fig. 2c appeared incorrectly. The corrected Fig. 2 is shown below.

Fig. 2 Tree density (a), mean DBH (b), skewness of DBH frequency distribution (c) and the number of recruits (d) of *P. glehnii* (solid circle) and *A. sachalinensis* (open circle)

The online version of the original article can be found at <http://dx.doi.org/10.1007/s11284-005-0093-2>

K. Takahashi
Department of Biology, Faculty of Science, Shinshu University,
Matsumoto 390-8621, Japan

M. Yogo
Department of Forest Science, Faculty of Agriculture,
Hokkaido University, Sapporo 060-0809, Japan

S. Ishibashi
Hokkaido Research Station, Forestry and Forest Products
Research Institute, Sapporo 062-8516, Japan

Present address: K. Takahashi (✉)
Department of Biology, McGill University,
1205 Docteur Penfield, Montreal,
Quebec H3A 1B1, Canada
E-mail: koichit@gipac.shinshu-u.ac.jp

Present address: M. Yogo
Tomoku Company Limited,
Marunouchi-Mitsui Building,
Marunouchi, Tokyo 100-0005, Japan